

From the Atherosclerosis Research Unit
Department of Medicine
Karolinska Institutet, Stockholm, Sweden

Peroxisome Proliferator-Activated Receptor Delta (PPARD)

Molecular studies of
regulation and activation

Petra Thulin



**Karolinska
Institutet**

Stockholm 2011

All previously published papers were reproduced with permission from the publisher.

The figure on the front page was adapted from Xu et al,
Proc Natl Acad Sci U S A. 2001 Nov 20;98(24):13919-24.

Published by Karolinska Institutet. Printed by Larerics Digital Print AB, Sweden

© Petra Thulin, 2011

ISBN 978-91-7457-203-2

To my family

ABSTRACT

The peroxisome proliferator-activated receptors (PPARs) are ligand-activated transcription factors involved in energy homeostasis. Their natural ligands are fatty acids and there are three different PPAR isoforms; PPARG, PPARG and PPARD. They are encoded by separate genes and have distinct functions, due to different tissue expression and affinity for ligands. PPARG controls genes involved in fatty acid oxidation, PPARG regulates genes important for fatty acid storage, and PPARD controls genes implicated in lipid oxidation and lipoprotein metabolism. Primates and humans treated with a PPARD agonist (GW501516) resulted in improved insulin sensitivity, increased HDL and decreased LDL cholesterol levels, making it a putative drug candidate for treatment of metabolic disease. PPARD has recently been assigned a beneficial role in macrophages, by inducing a switch from proinflammatory (M1) to antiinflammatory (M2) macrophages.

To characterize additional target genes of PPARD involved in the lipoprotein metabolism, the effect of PPARD activation on the apolipoprotein A-II (apoA-II) gene was investigated in human hepatoma cells. ApoA-II is one of the major proteins in the HDL particles. Treatment with GW501516 increased apoA-II promoter activity and mRNA levels in hepatoma cell lines. A site located at -737/-717 in the promoter was identified as the functional PPAR response element (PPRE). These results suggest that increased expression of the apoA-II gene is one of the reasons for the beneficial effects on lipoprotein metabolism after treatment with the PPARD agonist.

To investigate whether PPARs could regulate the alanine aminotransferase (ALT) genes, the effect of PPARG, G and D agonist treatment was studied. ALT activity in plasma is used as a marker for hepatotoxicity in humans. During a clinical trial with the PPARG ligand, AZD4619, the plasma ALT activity increased in some patients and *in vitro* studies showed that ALT1 protein and mRNA expression was induced by treatment with PPARG agonists in primary hepatocytes. Similarly, transient transfection of a promoter construct of ALT1 in HuH-7 cells showed increased activity mediated via a PPRE located at -574 after treatment with PPAR agonists. This study shows that the ALT1 gene is regulated by PPARs and that PPAR drugs might contribute to increased ALT activity in serum.

To explore regulation of the PPARD gene by posttranscriptional events, 5'- and 3'-RACE were performed on cDNA obtained from placenta, adipose tissue and pancreas. Both 5'- and 3'-alternative splicing of PPARD was identified. Coupled transcription/translation showed that the length and number of upstream AUGs in the 5'-UTR had a major impact on translational efficiency. Further, the promoter located upstream of exon one was verified as the major promoter, using reporter gene assays. A 3'-splice variant encoding a truncated PPARD protein, PPARD2, was shown to be a negative regulator of the full length receptor, PPARD1, in transient transfection assays.

To identify whether PPARD is regulated by microRNA (miRNA), the 3'-UTR was analysed *in silico*. Two putative miRNA target sites were identified in the PPARD 3'-UTR; miR-9 and miR-29. The miR-9 was verified as a functional miRNA targeting PPARD. However, PPARD mRNA levels remained unaffected by miR-9 expression, indicating that only the translation of PPARD was inhibited. Since both miR-9 and PPARD have been shown to play important roles in the inflammatory response of monocytes, the regulation of PPAR expression by miR-9 was investigated in these cells. A suppressive role of miR-9 on PPARD expression was identified in monocytes after LPS treatment but not in M1 or M2 macrophages, suggesting that the regulatory role of miR-9 on PPARD is exerted in monocytes, before differentiating into macrophages.

In summary, this thesis describes additional functions and ways of regulation of the ubiquitously expressed transcription factor PPARD with a major role in both health and disease.

POPULÄRVETENSKAPLIG SAMMANFATTNING

Hjärt- och kärlsjukdomar såsom hjärtinfarkt, hjärtsvikt och stroke (slaganfall) är de vanligaste dödsorsakerna i västvärlden. Hjärt- och kärlsjukdomar beror av en fettansamling i kärlen under många decennier som med tiden leder till åderförkalkning, så kallad ateroskleros. När åderförkalkningen blockerar kärl drabbas vävnader av syrebrist vilket leder till sjukdom. Uppkomst av åderförkalkning beror av flera faktorer, där övervikt, höga blodfetter, högt blodtryck och högt blodsocker är de viktigaste. Dessa faktorer påverkas i sin tur av ärftlighet men även av våra livsvanor, där rökning, fysisk inaktivitet, samt matvanor spelar en huvudsaklig roll.

Peroxisom proliferator-aktiverade receptorer (PPAR) är en grupp proteiner som reglerar gener som håller kroppen i jämvikt vid ökat eller minskat energiintag. De fungerar genom att binda till styrregioner i gener med betydelse för inlagring eller förbränning av fett. Det finns tre olika medlemmar i PPAR-familjen; PPAR alfa (PPARA), PPAR gamma (PPARG) och PPARD delta (PPARD). PPARA finns mest i levern, PPARG i fettväv medan PPARD finns i alla kroppens celler. Det har visat sig att aktivering av PPARD med en syntetisk molekyl, "GW501516" ökar mängden av det "goda" HDL-kolesterolet och minskar det "onda" LDL-kolesterolet. Syftet med denna avhandling var att identifiera gener som styrs av PPARD, samt studera hur nivåerna av PPARD styrs.

I den första studien undersöks resultatet av aktivering av PPARD i leverceller med GW501516 på apolipoprotein A-II (apoA-II) -genen. ApoA-II är ett vanligt förekommande protein i HDL-partikeln. Studien visar att PPARD kan binda till ett regulatoriskt område i apoA-II-genen och öka dess aktivitet, vilket skulle kunna tyda på att det är en av mekanismerna för hur PPARD-aktivering kan öka mängden HDL-kolesterol.

I den andra studien undersöks hur en potentiell läkemedelskandidat som binder PPARA, AZD4619, påverkar alaninaminotransferas (ALT), ett leverenzym som traditionellt används för att mäta leverskada. I en klinisk studie med AZD4619 observerades höjningar i plasmanivåerna av ALT, tydande på leverskada. Vår studie visar att aktivering av PPARA men även av PPARG och PPARD, kan styra ALT-genen och på så sätt öka mängden ALT-protein i cellerna. Slutsatsen är att ALT-höjningarna i den kliniska studien med AZD4619 inte nödvändigtvis berodde på leverskada utan kan ha orsakats av en regulatorisk effekt av PPARA.

Den tredje studien visar att nivåerna av PPARD kan styras genom alternativ splicing. För att en gen ska kunna ge upphov till ett protein krävs ett mellansteg där messenger-RNA (mRNA) bildas. För PPARD kunde vi se att det fanns olika typer av mRNA där längre varianter inte gav upphov till protein lika lätt som kortare varianter. Vi fann också en kortare PPARD proteinvariant, PPARD2, som kunde hämma funktionen av fullängdsvarianten, PPARD1. Slutsatsen är att alternativ splicing reglerar mängden av tillgängligt PPARD protein i kroppen, vilket skulle kunna ha betydelse vid exempelvis förändringar beroende på fysiologiska omständigheter i olika organ.

Den fjärde studien visar att PPARD kan styras genom regulatoriska molekyler som kallas micro-RNA (miRNA) som binder till mRNA-molekyler av PPARD och hämmar dem från att bli protein. Studien visade att miR-9 men inte miR-29 kunde hämma PPARD. Vid behandling med inflammatoriska substanser, då mängden miR-9 ökar, kunde vi se att effekten av PPARD på sina målgener minskade i monocytter, som är vita blodkroppar ansvariga för en snabb effekt av immunförsvaret vid en infektion. Vi undersökte även om det var skillnad i miR-9-mängd mellan proinflammatoriska M1- och antiinflammatoriska M2 makrofager, men såg ingen skillnad mellan dessa subtyper. Slutsatsen är att miR-9 kan ha en effekt på PPARD i monocytter men verkar ha mindre betydelse i makrofager.

Studierna i denna avhandling har visat att apoA-II- och ALT1-generna styrs av PPARD samt att mängden PPARD kan moduleras genom de regulatoriska mekanismerna alternativ splicing och miRNA-inhibering.

LIST OF PUBLICATIONS

This thesis is based on the following original paper, which are referred to by their Roman numerals:

- I. **Thulin P**, Glinghammar B, Skogsberg J, Lundell K and Ehrenborg E.
PPAR delta increases expression of the human apolipoprotein A-II gene in human liver cells
International Journal of Molecular Medicine 2008;21(6):819-24
- II. **Thulin P**, Rafter I, Stockling K, Tomkiewicz C, Norjavaara E, Aggerbeck M, Hellmold H, Ehrenborg E, Andersson U, Cotgreave I and Glinghammar B.
PPAR alpha regulates the hepatotoxic biomarker alanine aminotransferase (ALT1) gene expression in human hepatocytes
Toxicology and Applied Pharmacology 2008;231(1):1-9
- III. Lundell K*, **Thulin P***, Hamsten A and Ehrenborg E
Alternative splicing of human peroxisome proliferator-activated receptor delta (PPAR delta): effects on translation efficiency and trans-activation ability
BMC Molecular Biology 2007;8:70
* Equal contributors
- IV. **Thulin P**, Werngren O, Cheung L, Fisher R, Grandér D, Corcoran M and Ehrenborg E
MicroRNA-9 regulates expression of peroxisome proliferator-activated receptor delta (PPARD) in human monocytes
Manuscript

CONTENTS

1	INTRODUCTION	1
1.1	Cardiovascular disease	1
1.1.1	Lipoprotein metabolism	2
1.1.2	Atherosclerosis	3
1.1.2.1	Macrophages in atherosclerosis	5
1.1.3	Drugs in the treatment of CVD	5
1.1.3.1	Drugs in the treatment of dyslipidaemia	5
1.1.3.2	Drugs in the treatment of type-2 diabetes	6
1.1.3.3	Hepatotoxicity	7
1.2	Peroxisome proliferator-activated receptors	8
1.2.1	Structure	9
1.2.2	Cofactors	9
1.2.3	Function	10
1.2.4	PPARA	11
1.2.5	PPARG	12
1.2.6	PPARD	12
1.2.7	Ligands	13
1.2.6.1	Endogenous ligands	13
1.2.6.2	Synthetic ligands	14

1.3 PPAR regulation	15
1.3.1 Transcriptional regulation	16
1.3.2 Post-transcriptional regulation	16
1.3.2.1 Alternative splicing	16
1.3.2.2 MicroRNA	18
1.3.3 Post-translational regulation	20
1.3.3.1 Phosphorylation.....	20
1.3.3.2 Ubiquitination and sumoylation.....	20
2 HYPOTHESIS AND AIMS	22
3 MATERIALS AND METHODS	23
4 RESULTS AND DISCUSSION.....	29
5 GENERAL DISCUSSION	40
6 CONCLUSIONS	43
7 ACKNOWLEDGEMENTS.....	44
8 REFERENCES	48
9 PAPERS 1-IV	59

LIST OF ABBREVIATIONS

ABCA1	ATP-binding cassette transporter A1
ACO	acyl CoA oxidase
AF	activation function
ALT	alanine aminotransferase
ApoA-I	apolipoprotein A-I
ApoA-II	apolipoprotein A-II
BCL-6	B-cell lymphoma 6 protein
BMI	body mass index
ChIP	chromatin immunoprecipitation
CM	chylomicron
CE	cholesteryl ester
CVD	cardiovascular disease
DBD	DNA-binding domain
DR	direct repeat
EMSA	electrophoretic mobility shift assay
FABP	fatty acid binding protein
FACS	fluorescence activated cell sorting
HDL	high density cholesterol
IFN γ	interferon gamma
IL	interleukin
LBD	ligand-binding domain
LDL	low density lipoprotein
LPL	lipoprotein lipase
LPS	lipopolysackaride
M-CSF	macrophage colony-stimulating factor
miR	microRNA
MMP	matrix metalloproteinase
NCoR	nuclear receptor corepressor
NR	nuclear receptor
PCR	polymerase chain reaction
PPAR	peroxisome proliferator-activated receptor
PPRE	PPAR response element
PUFA	polyunsaturated fatty acid
RACE	rapid amplification of cDNA ends
RISC	RNA-induced silencing complex
ROS	reactive oxygen species
RXR	retinoid X receptor
SMART	silencing mediator for retinoic acid and thyroid hormone receptor
SPPARM	selective PPAR modulators
SUMO	small ubiquitin-like modifiers
TG	triglyceride
TNF α	tumour necrosis factor alpha
ULN	upper level of normal
UTR	untranslated region
VLDL	very low density lipoprotein

PREFACE

In this thesis I will introduce the research field of cardiovascular disease and the importance of the regulation and activation of the nuclear receptor PPARG. However, before doing so I would like to explain my reasons for choosing this particular topic. Ever since I was a child I have enjoyed running and jumping, and for many years I participated in track and field events. Naturally I became fascinated by physiology and the human body, so after graduating from Upper Secondary School, I studied “Integrative Human Medicine” for one year at Karolinska Institutet. In this time the topic of transcription factors was introduced to me and during a ten-week literature study about the AhR nuclear translocator (ARNT), my interest for research and transcription factors grew. A few years later, during my Master’s in Biomedicine I was introduced to the nuclear receptor PPARG in a lecture given by Associate Professor Ewa Ehrenborg. I was fascinated by the concept of a transcription factor that had important functions in metabolism but that could also be modulated by exercise. I was very happy when Ewa accepted me as a summer student in her group and after finishing my Master’s thesis on PPARG I was offered a PhD position and could not resist the opportunity.

1 INTRODUCTION

1.1 CARDIOVASCULAR DISEASE

Cardiovascular diseases (CVD) include a group of disorders of the heart and blood vessels which constitutes the major cause of death in the Western world today [1]. Myocardial infarction (MI) and stroke are the main killers of this group, although peripheral vascular disease also plays a role. The main risk factors for CVD are genetic predisposition and smoking together with the components of the metabolic syndrome, which include hypertension, dyslipidaemia, obesity (abdominal) and decreased glucose tolerance (i.e. insulin resistance) (Fig. 1) [2]. The world-wide increase in obesity is a major contributor to the increasing prevalence of the metabolic syndrome and the changes in our life-styles that have occurred during the last decade are to blame [3]. The access to high-caloric food and drinks has increased in combination with sedentary occupations and leisure. In the US about 68 % of the population is reported to be overweight (BMI >25) and 34 % are obese (BMI >30) compared to 45 % and 13.5 % fifty years earlier. The corresponding numbers in Sweden today is 45 % overweight and 12 % obese, respectively [4]. Accordingly, the numbers of people developing type-2 diabetes are increasing throughout the world and these individuals are at higher risk of developing CVD. The fact that the prevalence of obesity and type-2 diabetes is increasing in children and adolescents is also alarming, both from a socioeconomic and public health point of view [5].




Figure 1. The central role of the metabolic syndrome in vascular disease

Multiple environmental and genetic factors contribute to the formation of the metabolic syndrome marked by hypertension, dyslipidemia, obesity and insulin resistance. These risk factors in turn contribute to initiation and progression of type-2 diabetes and the macrovascular diseases, i.e. myocardial infarction, stroke, and peripheral vascular disease. (Adapted from Razani et al. 2008)

1.1.1 Lipoprotein metabolism

To understand the mechanisms for development of CVD, a brief introduction to the metabolism of lipids in the body is necessary. Dietary lipids are transported between the tissues of the body in the bloodstream as lipoprotein particles (Fig. 2) [6], which consist of a core of lipid surrounded by a shell of phospholipids and apolipoprotein, proteins specific for the destination of the lipoprotein particles. The lipoprotein particles include the chylomicrons (CM), very low density lipoproteins (VLDL), low density lipoproteins (LDL) and high density lipoproteins (HDL) and they all have specific compositions of lipids and apolipoproteins [7]. The lipoprotein metabolism is involved in determining the concentration of lipids in plasma and thus influences the amount of fat accumulation in tissues and arteries over a life span. Imbalance between production and removal of plasma lipids affects the homeostasis of the system and results in diseases caused by arterial lipid accumulation, such as atherosclerosis. Genes encoding for transcription factors that regulate proteins involved in the lipoprotein metabolism are central for the homeostasis of this system [8].




Figure 2. Schematic overview of the lipoprotein metabolism

Simplified overview of the lipoprotein metabolism showing the major lipoproteins and their routes. See text for details. (Adapted from Enkhmaa et al. 2010)

Dietary lipids are degraded by pancreatic enzyme, subsequently absorbed by the intestinal mucosa cells and packed into the CMs. The CMs mainly contain triglycerides (TG) and some cholesteryl esters (CE) in the lipid core, and the apolipoprotein B48 in the outer shell. The CMs are released into the blood stream and the TGs in the particles are hydrolyzed into free fatty acids by the enzyme lipoprotein lipase (LPL), which is

located on the surface of the endothelium in the tissues. The fatty acids are taken up primarily by skeletal muscle and adipose tissue and the CM remnants, which are enriched in CEs, are recycled by the liver. The liver produces and secretes VLDL particles, which consist primarily of TGs but also CEs, and they are labeled with the apolipoprotein B100. The function of VLDL is to provide the peripheral tissues with lipids, just like the CMs. As the VLDLs move through the circulation, they are depleted of triglycerides and enriched in cholesterol and become so called LDL particles. The LDL particles provide the tissues with cholesterol and are considered to be atherogenic. On the contrary, the HDL particles are responsible for the reverse cholesterol transport which means that HDL delivers cholesterol back to the liver, where it can be further metabolized and excreted as bile. The HDL particles are synthesized by the liver and they mainly contain the apolipoproteins A-I and A-II. These particles can interact with cholesterol transporters on the surface of cells in the periphery; ABCA1 in macrophages and ABCG1 in endothelial cells, which loads them with cholesterol. As the HDL particles return to the liver they are captured by the scavenger receptor B, class 1 receptors (SRB1) [9]. The antiatherogenic function of apoA-I is established, whereas the role of apoA-II has been more debated [10]. The effects of PPARD activation on apoA-II expression will be investigated in paper I.

1.1.2 Atherosclerosis

Atherosclerosis is the underlying reason for development of CVD. It is a process resulting in the formation of a lipid-rich plaque in large and medium-sized elastic and muscular arteries, which in case of rupture causes sudden thrombotic occlusion of the artery at the site of disruption (Fig. 3) [11]. In the heart, atherosclerosis can lead to myocardial infarction, whereas in the arteries of the brain, it can cause ischaemic stroke. Furthermore, atherosclerotic lesions in other arterial branches, might result in renal impairment, hypertension and critical limb ischaemia [12, 13]. The pathophysiology of atherosclerosis is not totally understood, but it involves the formation of lesions in the arteries which are characterized by lipid accumulation, inflammation, cell death and fibrosis. The fatty streaks are the earliest signs of lesions and they are present in the arteries already early in life and with increased age these fatty streaks might develop into atherosclerotic plaques [14]. A major risk factor to develop CVD is dyslipidaemia, characterized by elevated triglyceride levels, high levels of circulating LDL cholesterol, and low levels of HDL cholesterol [15]. The LDL cholesterol, especially the small dense LDL particles [16], accumulate in the

intima of the arterial vessels where they are oxidized, which induces endothelial expression of adhesion molecules such as vascular cell adhesion molecule 1 (VCAM-1) [17, 18]. These molecules attract monocytes from the blood stream to enter the intimal part of the artery, where they differentiate into macrophages under the influence of macrophage colony-stimulating factor (M-CSF) produced by endothelial cells and smooth muscle cells [19], and start expressing scavenger receptors [20]. The monocytes and macrophages belong to the innate immunity, the first line of defense of the body, and the scavenger receptors on macrophages detect oxidized LDL and promote their phagocytosis. As the disease progresses, the macrophages are overloaded with cholesterol, and turn into foam cells that start to produce inflammatory molecules which initiate an inflammatory response of the adaptive immunity, recruiting T-cells into the plaque. Activated T-cells produce inflammatory cytokines including interferon γ ($\text{IFN}\gamma$) and tumour necrosis factor α ($\text{TNF}\alpha$), which leads to further activation of macrophages and endothelial cells [12]. The activated macrophages produce additional $\text{IFN}\gamma$ and $\text{TNF}\alpha$, as well as matrix-degrading enzymes, matrix metalloproteinases (MMPs), and reactive oxygen species (ROS). $\text{IFN}\gamma$ and $\text{TNF}\alpha$ inhibit smooth-muscle-cell proliferation and collagen production, which increases the vulnerability of the plaque and hence the risk of rupture [21]. The MMPs further destabilize the plaque by degradation of the protective collagen and elastin [22], whereas the ROS contribute to oxidative modification of LDL and oxidative damage of DNA [23]. Furthermore, $\text{TNF}\alpha$ inhibits the enzyme LPL, leading to hypertriglyceridemia, which also promotes atherogenesis [24].




Figure 3. Simplified illustration of the initiation and progression of the atherosclerotic plaque
The importance of macrophages in the atherosclerotic process is emphasized. See text for details.
(Adapted from www.invivo.ca/illustration_and_print_format_of_atherosclerosis.html)

1.1.2.1 Macrophages in atherosclerosis

Recently, different macrophage subsets have been identified in the atherosclerotic lesions, which are important for the inflammatory status of the plaque. The “classically activated” M1 macrophages are induced by LPS and Th1 cytokines, mainly IFN γ , and they produce inflammatory mediators, such as IL-6 and TNF α . In contrast, the “alternatively activated” M2 macrophages develop as a response to the Th2 cytokines IL-4 and IL-13 and they secrete antiinflammatory mediators, such as transforming growth factor beta (TGF β) and IL-10, to dampen the inflammatory response [25]. Both macrophage subpopulations are expressed in atherosclerotic lesions and modulate the inflammatory response, thereby they might have an impact on plaque stability and hence risk of development of CVD [26]. However, there is a plasticity between the subtypes and activated M1 macrophages can be modulated into M2 macrophages by expression of Th2 cytokines and vice versa [27]. It has been shown that expression levels of PPAR γ in atherosclerotic lesions are correlated with M2 markers, suggesting that PPAR γ plays a beneficial role in atherosclerosis by regulating macrophage subclasses. Also, PPAR γ activation primes human primary monocytes into the alternatively activated phenotype [26]. Accordingly, myeloid specific deletion of PPAR γ in mice impairs alternative activation and predisposes these animals to development of diet-induced obesity, insulin resistance, and glucose intolerance [28]. Additionally, the PPAR family member PPAR δ might be a major determinant of macrophage subtype activation as its expression is essential for alternative activation of resident macrophages in the liver and adipose tissue [29, 30]. Loss of PPAR δ expression in the macrophages in these tissues results in insulin resistance and obesity, respectively. However, a recent report did not show any correlation between PPAR δ mRNA expression and M2 markers in human atherosclerotic lesions [31] and this subject will be further investigated in paper IV.

1.1.3 Drugs in the treatment of CVD

1.1.3.1 Drugs in the treatment of dyslipidaemia

It has been concluded that the most important and cost-effective interventions for the atherogenic lipid phenotype are an appropriate diet, weight loss, exercise, and smoking cessation [32]. In addition, there are pharmacological agents that can improve the metabolic profiles in already affected individuals. Most of the drugs in use today have effects on lowering of LDL cholesterol, since it has been considered the most

important parameter of the atherogenic dyslipidaemia [33]. However, low plasma HDL cholesterol has also been recognized as an independent risk factor of CVD and should be treated in addition to the LDL cholesterol [34-36]. Statins are the drugs of first choice for patients with high plasma concentrations of cholesterol commonly observed in dyslipidaemia, CVD, type-2 diabetes and other high risk states of atherosclerosis. Statins are primarily LDL cholesterol lowering agents and inhibit hydroxymethylglutaryl-CoA reductase, which catalyses the rate-limiting step in the endogenous cholesterol synthesis [33]. However, there are compounds which can be used as alternative drugs if statins are not well tolerated or in combination with statins if the desired effect is not achieved. The fibrates are drugs that activate PPARA in muscle, liver, and other tissues. They mainly decrease hepatic VLDL secretion and therefore plasma triglycerides, but fibrates also decrease the levels of LDL cholesterol in individuals with elevated baseline plasma LDL concentrations and increase HDL cholesterol levels in patients with low baseline plasma concentrations of HDL [37, 38]. Nicotinic acid (also known as Niacin) decreases both LDL cholesterol and triglycerides and it is the strongest HDL raising drug in clinical use at the moment. However, the main limitation for the use of nicotinic acid is its low tolerability, as it causes flush in almost every patient [33]. In addition the anion exchange resins, cholesterol absorption inhibitors and omega-3-fatty-acids are all compounds that can be used as combination therapies together with statins, however the effects on cardiovascular end-points are not established for the two latter ones. Large ongoing trials address the decisive question whether treatment with fibrates and niacin provides additional cardiovascular risk reduction when given in addition to statin treatment [33].

1.1.3.2 Drugs in the treatment of type-2 diabetes


In order to control glucose levels in type-2 diabetes patients, the first line of drugs have been metformin and sulfonylureas. Sulfonylureas increase the secretion of insulin from the pancreatic beta cells [39] whereas metformin has been identified as an activator of AMP kinase, reducing glucose output from the liver and increasing the peripheral glucose utilization, in addition to its direct vascular effects [40]. The thiazolidinediones (TZDs) or glitazones are PPARG agonists which increase insulin sensitivity due to the uptake of free fatty acids from the circulation by the adipose tissue [41]. In addition there are some new insulin sensitizing drugs out on the market, which will not be discussed in this thesis.

1.1.3.3 Hepatotoxicity

One of the most common reasons for stopping the development of promising drug candidates is hepatotoxicity discovered during preclinical studies. Furthermore, nearly 50% of all drugs fail in the post-marketing phase due to unexpected toxicity or unwanted interference with metabolism issues [42]. Since the introduction into clinical monitoring some 50 years ago, serum alanine aminotransferase (ALT) has become the standard biomarker for detection of liver injury [43]. ALT activity has been detected mostly in liver tissue [44] and ALT in serum is believed to be due to leakage from damaged hepatocytes. Serum aspartate aminotransferase (AST) activity is considered a less specific biomarker of liver function compared to ALT, since AST is present to a higher degree in skeletal and cardiac myocytes than in hepatocytes [45]. Elevations in ALT activity show high correlations to liver damage caused by drug toxicity, infection, alcohol, cirrhosis and inflammatory steatosis, and ALT activity is usually between 10-100 times higher than normal levels in these states [46]. However, the function of ALT as a perfect liver injury marker has recently been questioned [47]. While it is true that the liver is the most ALT rich organ, ALT is also found at high concentrations in kidney, cardiac and skeletal muscle [48]. Furthermore, ALT is an important enzyme for gluconeogenesis and amino acid metabolism in humans (Fig. 4) [49]. Approximately 10% of the patients taking fenofibrate show increased aminotransaminases above normal levels. Despite this, there has been no reports of hepatic pathologies with fenofibrate use [50, 51] and in 1998, Edgar et al suggested that fibrates could increase the gene expression of ALT as an alternative “non-toxic” mechanism for the elevation of ALT enzymes in serum [46]. The normal turnover of hepatocytes, releasing their increased amounts of intracellular aminotransferases, would be the reason for the mild increase in serum ALT observed after treatment with fibrates. The subject of a possible PPAR-mediated regulation of the ALT1 gene will be investigated in paper II in this thesis.

Figure 4. Schematic overview of the glucose-alanine cycle catalysed by the enzyme ALT

ALT is responsible for transamination of amino acids in the muscle and for gluconeogenesis in the liver. The amino groups transported from the muscle to the liver are converted into urea which is excreted in the urine. (Adapted from www.themedicalbiochemistrypage.org)



1.2 PEROXISOME PROLIFERATOR-ACTIVATED RECEPTORS

Peroxisome proliferator-activated receptors (PPARs) belong to the superfamily of nuclear receptors (NRs) which contain transcription factors that regulate the expressions of genes involved in processes such as reproduction, development and general metabolism. There are 48 members of the NR superfamily in humans and 49 in the mouse [52] and they all share a common structure. The PPARs are classified as members of group C in the nuclear receptor subfamily 1 (NR1C) [53] and they belong to a functional subgroup of the nuclear receptors denoted adopted orphan receptors (Fig. 5) [54].




Figure 5. Functional grouping of the nuclear receptor superfamily

The PPAR subfamily belongs to the group of adopted orphan receptors and is indicated in the figure with an arrow. (Adapted from Chawla et al 2001)

PPARA (NR1C1) was the first member of the PPAR family to be discovered in the early 1990s [55] and obtained its name from the fact that it could bind agents that caused peroxisome proliferation in rodents. PPARB (NR1C2) and PPARG (NR1C3) were identified a few years later in *Xenopus* [56] and subsequently murine PPARG and one more paralogue which was named PPARD were identified [57]. However, when the chicken PPARB was cloned and characterized, it was evident that murine PPARD was the ortholog of *xenopus* PPARB and hence this receptor is now referred to as PPARB/D. In this thesis it will exclusively be referred to as PPARD. The PPARs are

expressed by separate genes on different chromosomes and they are expressed in a tissue-specific manner.

1.2.1 Structure

All the nuclear receptors share a common structure which makes them able to bind to DNA and exert their functions as regulators of gene transcription (Fig. 6) [54]. The N-terminal ligand-independent activating function 1 (AF-1) domain is the most variable between the PPAR family members, and it is recognized by coactivators and is considered to be important for transactivation and transcription [58]. The DNA-binding domain (DBD) is very well conserved and it is constituted by two highly conserved zinc finger-like structures which are responsible for receptor binding to its specific DNA sequences, the PPAR response elements (PPRE), and it also plays a role in dimerization [59]. The hinge region (H) permits protein flexibility to allow simultaneous receptor dimerization and DNA binding [54]. The ligand binding domain (LBD) consists of 12 α -helices (H1-12) which together constitute a pocket for the ligand. The entrance to the pocket is guarded by H12, which contains residues that are crucial for the function of AF2. In the absence of ligand, corepressors are bound to H3 and H4 but when a ligand is bound, H12 induces a conformational change in the receptor and the corepressors are released. Instead, coactivators which contain LxxLL-like motifs (where L is leucine and x is any amino acid) can bind to a hydrophobic cleft formed by H3, H4 and H12 which supports coactivator interaction and gene transcription. The LBD is also essential for dimerization. [60]



Figure 6. Schematic representation of the functional domains of PPAR

The N-terminal domain contains the activation function 1 (AF-1). DBD = DNA-binding domain, H = hinge region, LBD = ligand-binding domain. Amino acid (aa) numbers are shown above the receptor. (Adapted from Ehrenborg and Krook 2009)

1.2.2 Cofactors

The nuclear receptors need additional proteins called cofactors to be able to exert their functions. The nuclear receptor cofactors include corepressors, coactivators and their associated proteins, and over 300 of them have been identified [61]. The coactivators directly bind to transcription factors and positively regulate target gene

transcription, either by modification of histone and chromatin structure to open up DNA for transcription, while others provide linkage to the core basal transcriptional machinery. CBP/P300 and SRC-1 are examples of coactivators which have been shown to associate with the PPARs and they loosen up the chromatin structure to enable transcription by possessing histone acetylase transferase (HAT) activity [62-65]. The corepressors inhibit target gene transcription by recruitment of histone deacetylases (HDACs) to enforce a tight chromatin structure and classical examples are NCoR and SMRT [66, 67]. As far as is known, there are no receptor specific coactivators that direct the overall transcriptional activity of particular members of the PPAR subfamily [68]. In the absence of ligand, the PPARs are maintained in the nucleus in a repressed state by nuclear receptor corepressors such as NCoR and SMRT [69-71]. PPARD is the only PPAR subtype which is able to bind DNA when bound to corepressors and it can thereby repress PPAR target genes in the absence of ligand [69, 70, 72]. Another mechanism of gene repression exerted by PPARD is its association with BCL-6, which functions as a corepressor for genes involved in inflammation [73]. In the unliganded state, PPARD binds to and sequesters B-cell lymphoma 6 protein (BCL-6), whereas ligand binding causes BCL-6 dissociation and repression of inflammatory genes. Binding of a ligand to a PPAR results in the dissociation of the corepressors and then an orchestrated recruitment of several transcriptional coactivators which facilitates gene transcription. Besides ligand binding, activation or repression of target genes by the PPARs also depends on the cellular expression pattern of coactivators and corepressors, as has been taken advantage of in the SPPARM concept (described in section 1.2.6.2) [74].

1.2.3 Function

The PPARs are lipid sensors with distinct but overlapping functions in glucose and lipid metabolism. The natural ligands for all the PPARs are unsaturated fatty acids and their derivatives. Upon ligand binding the PPARs heterodimerize with the retinoid X receptor (RXR) and bind to PPRE which consists of a direct repeat (DR) of the consensus sequence AGGTCA separated by one nucleotide, called a DR1 motif [75] (Fig. 7) [58]. It has been reported that the 5'-flanking region to the PPRE is responsible for subtype specific activation of the PPARs [76]. The conformational change induced by ligand activation releases corepressors and allows coactivators to bind, which induces transactivation and transcription.




Figure 7. PPAR ligand binding, dimerization and transactivation

Upon ligand binding the PPARs form heterodimers with RXR and bind to PPARE in gene promoters. The effects are further modified by interaction with specific cofactors. Fatty acid binding proteins (FABP) may bind and deliver ligands to the receptor. (Adapted from Ehrenborg and Krook, 2009)

1.2.4 PPARA

The human *PPARA* gene consists of eight exons and it is localized at chromosomal region 22q12-q13.1 [77, 78]. PPARA is primarily expressed in tissues with a high level of fatty acid metabolism such as liver, skeletal muscle, heart, kidney, and intestine [79]. The major role of PPARA is the regulation of energy homeostasis and mainly in the liver, PPARA activates fatty acid catabolism, stimulates gluconeogenesis and ketone body synthesis [80, 81]. The function of PPARA in hepatic fatty acid metabolism is especially prominent during fasting and in normal mice, PPARA expression increases during fasting. In contrast, fasted PPARA knock-out mice have fatty livers and elevated levels of free fatty acids in plasma due to the inability of the liver to oxidize the fatty acids released from adipose tissue [81]. Furthermore, fasted PPARA null mice show decreased levels of plasma glucose and ketone bodies due to decreased gluconeogenesis and lipid oxidation, respectively. In addition, the starved PPARA mice suffer from hypothermia [81]. An additional function of PPARA is to control the hepatic expression of enzymes involved in the control of amino acid metabolism and urea synthesis [82]. PPARA is also involved in the regulation of lipoprotein assembly in the liver, which is demonstrated by the capacity of the PPARA activating fibrates to increase the expression of apolipoproteins involved in the reverse cholesterol transport [83-85]. PPARA has also been described to be a modulator of inflammation by negative crosstalk with the NF κ B pathway and treatment of hyperlipidemic patients with fibrates is shown to decrease the elevated levels of TNF α and IFN γ [86-88].

1.2.5 PPARG

The human *PPARG* has nine exons and is located on chromosome 3p25 [89]. The human *PPARG* gene gives rise to three different transcripts due to alternative promoter usage (*PPARG*1-3) but after alternative splicing of the 5' -end of the transcripts, *PPARG*1 and 3 gives rise to the same protein, hence there are two protein isoforms of the human *PPARG* (*PPARG*1 and *PPARG*2) [90]. The functional difference between the isoforms is not clearly understood but *PPARG*2 expression is restricted to adipose tissue where it seems to play a major role in adipocyte differentiation, whereas *PPARG*1 is more widely expressed. Adipose tissue, large intestine, and hematopoietic cells express the highest amounts of *PPARG*1 while kidney, liver, and small intestine have intermediate levels. *PPARG*3 is abundant in macrophages, the large intestine and white adipose tissue [90]. The function of *PPARG* is to induce adipocyte differentiation and increase the expression of genes involved in lipogenesis [91]. Furthermore, *PPARG* is involved in the glucose metabolism through an improvement of insulin sensitivity, when fatty acids are taken up from the circulation. Lately *PPARG* has also been recognized to play an important role in inflammation, favouring a more antiinflammatory phenotype of the residential macrophages in adipose tissue and liver [28].

1.2.6 PPARD

Human *PPARD* is located at chromosomal region 6p21.2-p21.1 and comprises nine exons [92]. *PPARD* is ubiquitously expressed and its function was an enigma for a long time. It is the least studied of the three *PPAR* subtypes due to its ubiquitous expression and unavailability of selective ligands [93]. However, during the last decade it has received more attention and its role in energy homeostasis and fatty acid catabolism has been recognized since the specific ligand GW501516 was synthesized. Administration of GW501516 to mice increases oxidation of fatty acids in the skeletal muscle and ameliorates diet-induced obesity and insulin resistance [94]. In insulin resistant rhesus monkeys, treatment with GW501516 showed decreased plasma triglyceride and LDL cholesterol levels, especially the small dense LDL particles, as well as increased plasma HDL levels. Additionally, the insulin levels of the monkeys were decreased by treatment with the *PPARD* agonist [95]. The importance of *PPARD* for lipid metabolism has also been demonstrated by transgenic tissue-specific expression in mice. Transgenic expression of an activated form of *PPARD* in adipose tissue results in lean mice which are resistant to obesity induced by a high-fat diet

[96]. In parallel, muscle-specific PPARD overexpression results in a shift to more oxidative muscle fibers (type 1), increased expression of genes involved in oxidative metabolism and reduced body fat mass [97]. Also, endurance exercise promotes an accumulation of PPARD protein in muscle of wild-type mice and transgenic mice overexpressing the activated form of PPARD in skeletal muscle, are lean and show increased endurance and expression of oxidative fibers compared to their wild-type littermates [97, 98]. The beneficial role of PPARD on lipid metabolism makes it a plausible target for treatment of obesity and insulin resistance. Studies in mice also reveal the importance of PPARD in the embryonic development, in the skin, brain and in the regulation of inflammatory macrophages in liver and adipose tissue [29, 30, 99, 100].

1.2.7 Ligands

1.2.7.1 Endogenous ligands

All the PPARs have wide ligand-binding pockets compared to the other members of the nuclear receptor family, which makes them able to bind fatty acids and their derivatives. Hence, the natural ligands for all the PPARs consist of polyunsaturated fatty acids (PUFAs), such as linoleic, linolenic and arachidonic acid [101]. However, there are some differences in ligand preference between them; the ligand binding cavity in PPARG is more hydrophobic than the other two PPARs, which explains why it also can accommodate saturated fatty acids. In contrast, PPARD is shown to have a narrower pocket compared to PPARG and PPARG, which explains why it is more restrictive in binding saturated fatty acids [102]. Furthermore, some eicosanoids can serve as subtype specific ligands, for example leukotriene B₄ can activate PPARG whereas 15-Deoxy-D12,14-PGJ₂ has been identified as a more specific ligand for PPARG [101]. In contrast, the fatty acid erucic acid (C22:1), which is a weak ligand, appears more selective for PPARD [103]. In addition to ligand specificity, there are intracellular lipid binding proteins, so called fatty acid binding proteins (FABP) which bind ligands specific for the different PPAR isoforms and bring them into the nucleus of the cell where the PPARs are located. Interestingly, although FABP4 and FABP5 bind multiple ligands, only particular compounds trigger their nuclear translocation [104]. FABP4 moves into the nucleus in response to ligands that activate PPARG but not upon treatment with PPARD ligands. In contrast, FABP5 mobilizes to the nucleus only in response to ligands that activate PPARD [105].

1.2.7.2 *Synthetic ligands*

Because of their central role in the regulation of genes involved in energy homeostasis and inflammation, PPARs have become attractive pharmaceutical targets for the treatment of metabolic disease. The fibrates activate PPARA and the use of fibrates in the management of lipoprotein disorders has a history dating back to the mid-1960s [106]. They show few significant side-effects and are considered to be relatively safe, but have been associated with muscle weakness, myopathy and in some rare cases rhabdomyolysis [107]. The glitazones are synthetic ligands for PPARG and were introduced on the market at the end of the 1990s and they are associated with weight-gain, fluid-retention and an increased risk of bone fracture [108-110]. Since the withdrawal of Troglitazone in the year 2000 due to hepatotoxicity [111] and now the recent withdrawal of Rosiglitazone from the market in 2010 due to the increased risk of MI, stroke and overall mortality [112, 113], the safety concerns of PPARG agonists have been raised. Regarding PPARD specific agents, no compounds are in clinical use yet, but there are some small recent human studies that have evaluated the short-term effects of GW501516 [114, 115]. Both studies showed decreased triglyceride levels after two weeks treatment despite the fact that the participants were healthy and only moderately obese. Increased HDL-levels was only obtained in one of the studies [114] whereas decreased LDL cholesterol and increased insulin sensitivity was observed in the other one [115]. However, GlaxoSmithKline, the company in charge of GW501516, terminated a phase I trial with this compound in 2005 and it is no longer on their list of compounds in the pipeline [116]. Another synthetic PPARD agonist, MBX-8025, from the company Metabolex showed decreased triglyceride and LDL cholesterol and increased HDL cholesterol in an eight week long study including 173 obese or overweight participants [117]. Considering the side-effects of some PPARG agonists, which have been out on the market for years, there is already reluctance towards PPAR compounds and it is obvious that any drug acting via PPARs needs to demonstrate a favorable cardiovascular risk-benefit profile. Dual PPARA/G agonists and pan-PPARA/G/D agonists have been developed in hopes of achieving multiple therapeutic benefits but reducing the side-effects [118]. However, the safety is still an issue for these agents. Another type of second generation PPAR compound is the selective PPAR modulators (SPPARMs) which are designed to activate the PPARs in a tissue-selective manner [74]. Hence, only desirable effects of PPAR activation are

achieved whereas side-effects can be avoided. The SPPARMs bind to the PPAR and induce a conformational change in the LBD distinct from when a full agonist is binding, resulting in preferential binding of specific cofactors or corepressors. Thus, differential PPAR effects can be achieved depending on tissue expression of distinct sets of coactivators and corepressors. The SPPAR γ Ms show good efficacy and less side-effects in pre-clinical animal models of metabolic disease and it remains to see how well they perform in clinical trials [74].

1.3 PPAR REGULATION

So far, the main focus of PPAR research has been to unravel the effects of activation of PPARs whereas considerably less is known about the regulation of their expression [119]. Most research has been performed using mice models, which means that the regulatory mechanisms between humans and mice might deviate. However, keeping that in mind, the main mechanisms of regulation of the PPARs will be summarized below and an overview of these mechanisms is shown in Figure 8.




Figure 8. Schematic overview of the main mechanisms of PPAR regulation

The levels of regulation are depicted on the left hand side and the mechanisms of regulation are described on the right hand side of the figure. The figure is based on the PPARG gene, since it is the main focus of this thesis, even though the mechanisms are valid for all the PPAR family members. The transcriptional start of the PPAR gene is indicated with an arrow and +1.

1.3.1 Transcriptional regulation

Since there is not much data regarding the transcriptional regulation of the PPARs, the main focus of this chapter will be on the posttranscriptional regulation of the PPARs. However, it can be mentioned that a recent study identified differential binding of estrogen-related receptor (ERR) α and γ to a site encompassing a single nucleotide polymorphism (SNP) in the human PPARA promoter, which influences the transcriptional activity of the gene [120]. Furthermore, adipogenic hormones such as insulin and glucocorticoids induce PPARG transcription during early adipogenesis through the CCAAT enhancer-binding proteins [121]. Also, the SREBP1 and -2 directly control the expression of the human PPARG gene [122]. *In silico* analysis of the human PPARD promoter showed that it lacks a TATA box but is rich in potential Sp-1 binding elements, which is a typical feature for house-keeping genes, and it also contains two putative NF κ B elements [92].

1.3.2 Post-transcriptional regulation

1.3.2.1 Alternative splicing

Splicing of precursor mRNA (pre-mRNA) is a crucial regulatory step in the pathway of gene expression. The mature mRNA is formed as the introns of the pre-mRNA are removed and the exons are ligated together (Fig 9) [123]. Splicing occurs in organisms ranging from yeast to human and take place within the spliceosome, a large protein complex containing the small nuclear RNAs (snRNAs) U1, U2, U4, U5 and U6, and more than 100 core proteins [124, 125] (Fig 9B) [123].

Four signals located at the exon-intron boundaries have a well characterized role in helping directing the splicing machinery. In the 5'-end of the intron, the 5' splice site denoted by the nucleotides GU in the exon/intron boundary encompassed within a larger less conserved sequence. In the 3'-end of the intron, there is the branch point containing an adenosine, a polypyrimidine tract (PPT), followed by a terminal AG [126] (Fig. 9A). In addition, there are short conserved regions in the exons and introns which act as exonic or intronic splicing enhancers or silencers. Specific binding of splicing regulatory proteins called SR proteins to these regulatory regions assists in the positioning of the spliceosome on the appropriate splice sites [127, 128].

Alternative splicing, which includes different exons in the mRNA, results in the generation of alternative isoforms and is often tightly regulated in a cell-type- or developmental stage-specific manner [129]. There are several different types of alternative splicing events (Fig. 9C) of which exon skipping is the most common in higher vertebrates, followed by alternative 3'- and 5'-splice site (SS) selection, and intron retention [130]. Also, more complex events like mutually exclusive exons, alternative promoter usage and alternative polyadenylation give rise to alternatively spliced transcripts in vertebrates [131].




Figure 9. Schematic overview of the mechanisms of splicing

A) The four conserved signals that enable recognition of RNA by the spliceosome are shown above. B) The splicing machinery that performs the main steps in the splicing process is shown. C) Different types of alternative splicing events. SS = splice site; PPT = polypyrimidine tract. (Adapted from Keren *et al* 2010)

Dominant negative regulation by truncated receptor isoforms has been described for many of the members of the nuclear receptor superfamily, such as the glucocorticoid receptor, estrogen receptor and vitamin-D receptor [132-134]. Both the PPARA and PPARG genes have been shown to express truncated isoforms by alternative splicing, which act as dominant negative regulators of their respective full-length receptor [135, 136]. The truncated isoforms lack a part of the ligand-binding domain, which makes them unable to bind ligands, and they are expressed at low levels compared to the full-length variants. Alternative splicing for the human PPARD gene has not been studied, even though a truncated version has been reported in the NCBI database. In paper III, the putative regulation of PPARD by alternative splicing is investigated.

1.3.2.2 *MicroRNA*

MicroRNAs (miRNAs) are endogenous non-coding single-stranded RNAs approximately 22 nucleotides long that bind to their target mRNA and either induce mRNA degradation or inhibit protein translation [137]. They are expressed from animals to plants, are conserved between species and are implicated in diverse biological processes such as development and differentiation [138]. Furthermore, many of them have been reported to be aberrantly expressed in different forms of cancers [139]. In mammals, more than 50% of mRNAs are predicted to be the subject of miRNA-mediated control [140]. The miRNAs are initially transcribed in the nucleus by RNA polymerase II as long primary transcripts (pri-miRNAs) which are capped and polyadenylated, and cleaved into 60-80 nucleotide hairpin structures (pre-miRNAs) (Fig. 10) [141]. After the completion of the nuclear processing, the pre-miRNA is exported into the cytoplasm by the nuclear transporter protein Exportin 5 where it undergoes further processing by the Dicer, releasing ~22 nt miRNA duplexes containing ~2 nt overhangs at either end. The strand with the least stable pairing in the 5'-end (the guide strand) is selectively loaded into the RNA-induced silencing complex (RISC) which targets transcripts through either mRNA cleavage or translational repression [142, 143]. The other strand (passenger strand or miRNA*) is degraded.

The miRNAs:RISC complexes bind to and regulate their target mRNAs by partial complementary binding in the 3'- untranslated region (3'-UTR) of the mRNA. However, there are exceptions to this rule and some miRNAs have been reported to bind to the 5'-UTR or to the coding region of the mRNA [144-147]. Additional features of the target mRNA has shown to be of significance such as AU-rich elements and positioning of the sites within the 3'-UTR [148]. The most important part of binding is between nucleotide 2-8 from the 5'-end of the miRNA, the so called seed sequence [149, 150]. In addition, complementarity in the 3' end of the miRNA has proven to be of significance for increased stability of the binding and compensation of a weak seed region [150]. Another recently discovered group of miRNAs are the ones with centered pairing, which lack both perfect seed pairing and 3'-compensatory pairing and instead have 11–12 contiguous Watson-Crick pairs to the center of the miRNA [151].




Figure 10. Biogenesis and function of miRNA

The miRNAs are transcribed in the nucleus and delivered into the cytoplasm where they exert their effects on mRNA degradation or protein synthesis. See text for details.

(Adapted from Chang and Mendell, 2009)

Transcription factors are often subjected to miRNA regulation, which introduces an additional level for the cell to control its regulators. Of the human PPARs, both PPARA and PPARG have been shown to be regulated by miRNAs. PPARA protein expression is targeted by miR-519d in subcutaneous adipose, a miRNA which is shown to be highly expressed in severely obese compared to non-obese subjects [152]. MiR-519d was also shown to increase lipid accumulation during preadipocyte differentiation, indicating that it plays a role in obesity. PPARG is targeted by miR-27 and miR-130, which are downregulated during adipocyte differentiation [153-156]. No data are yet available regarding the regulation of PPARD by miRNA, however the question of whether PPARD is miRNA regulated will be investigated in paper IV.

1.3.3 Post-translational regulation of PPARs

1.3.3.1 Phosphorylation

Both mPPARA and mPPARG are shown to be phosphorylated by various stimuli affecting the different domains of the receptors. For example, insulin activates MAPK which activates ERK leading to phosphorylation of the A/B domain in AF-1 containing N-terminus of the receptor, resulting in release of corepressors and increase in the transactivational activity of both receptors [157]. So far, less information is available regarding the phosphorylation of the PPARD isotype, even though there are sites predicted in the protein, which implies that PPARD also might be a phosphoprotein [158]. Furthermore, both cAMP and PKA activators are shown to increase both ligand-activated as well as basal activity of PPARD [159, 160].

1.3.3.2 Ubiquitination and sumoylation

Ubiquitin is an 8 kDa protein which covalently binds to proteins to target them for degradation by the 26S proteasome. *In vitro* studies have shown that both PPARA and PPARG can be targeted for degradation by ubiquitination. Furthermore, PPARA agonists stabilize the receptor whereas the degradation of PPARG via ubiquitination is enhanced by ligand binding [161, 162]. For PPARD, *in vitro* studies show that under conditions of moderate expression, GW501516 is not significantly influencing ubiquitination or degradation of PPARD. In contrast, overexpression of PPARD via transient transfection dramatically enhances its degradation by ubiquitination and proteosomal degradation, which is inhibited upon ligand binding [163].

Small ubiquitin-like modifiers (sumo) compose a family of three 11 kDa proteins homologous to ubiquitin which can be reversibly conjugated to the lysine residues of proteins through covalent binding [164]. Many of the sumo-modified proteins identified to date are transcription factors, coactivators, or corepressors and in the majority of cases attachment of sumo appears to repress the activity of transcriptional activators [165]. So far, among the PPAR family members PPARA [166, 167] and PPARG [168] have been described to be regulated by sumoylation. Modification of PPARA by sumoylation specifically recruits the corepressor NCoR but not SMRT which might lead to differential expression of a subset of PPARA target genes. For PPARG, it is shown that ligand-dependent sumoylation directs the receptor to the promoter of the iNOS gene after inflammatory stimuli and induces its transrepression,

which explains the inhibitory effect of PPARG on some genes involved in inflammation [168].

It is obvious from this introduction that further studies regarding the regulation of the PPARs are needed, especially of PPARD. Furthermore, additional PPARD target genes implicated in metabolic disease remain to be identified. Clearly this receptor has its beneficial properties, even though its broad tissue expression might limit its potential as a drug target. Hence, this thesis will explore additional target genes of PPARD as well as further elucidate the mechanisms of its regulation.

2 HYPOTHESIS AND AIMS

Hypothesis:

The overall hypothesis for this thesis is that PPARD activation increases the expression of genes beneficial to prevent the development of atherosclerosis and CVD, and furthermore that the PPARD gene per se is subjected to regulation by posttranscriptional mechanisms.

Aims:

The general aims of this thesis were to identify human PPAR target genes and improve our understanding of the molecular mechanisms regulating the expression of human PPARD.

Specific objectives

- To characterize the mechanisms of PPARD activation on the apoA-II gene expression (paper I)
- To determine whether the ALT genes are transcriptional targets of the PPARs (paper II)
- To investigate whether PPARD is subjected to posttranscriptional regulation via alternative splicing and miRNA (paper III and IV)

3 MATERIALS AND METHODS

Detailed descriptions of materials and methods are given in the individual papers.

3.1.1 Chemicals (paper I-IV)

The synthetic PPAR δ agonist GW501516 was synthesized by Synthelec AB as described [169]. Fenofibric acid was a kind gift from Professor Per Eriksson, Karolinska Institutet, Sweden. Additional fenofibric acid and AZD4619 were obtained from Astra Zeneca, Mölndal, Sweden. Rosiglitazone was obtained from Cayman Chemical Company and TNF α from Calbiochem. The cytokines used in paper IV were purchased from Peprotech and LPS from Sigma.

3.1.2 Bioinformatic sequence analyses (paper I-IV)

In order to find putative PPREs in the hypothesized target genes apoA-II and ALT1, the MatInspector program in the Genomatix database [170], the Alibaba 2.1 [171] and Promo [172] programs were used (paper I and II). To identify sequence conservation between species the ECR browser tool [173], and ClustalW [174] were utilized (paper I and III). Furthermore, the RepeatMasker [175] was used to look for repetitive elements in the PPAR δ gene. In paper IV, miRNA target site prediction was performed by the UCSC genome browser [176] (human May 2004 assembly), which includes miRNA target predictions from the databases TargetScanS [177] and PicTar [178].

3.1.3 Cell culture (paper I-IV)

A panel of human cell lines have been used in this thesis; the human hepatocellular carcinoma cell lines HepG2 and HuH-7, the human embryonic kidney cell line HEK293, the cervical epithelioid carcinoma cell line HeLa, and the kidney epithelial carcinoma cell line A498 were all maintained in DMEM (1 g/l glucose), containing 10 % FBS, penicillin (100 U/ml) and streptomycin (100 μ g/ml) at 37°C in 5 % CO $_2$ in air. The monocytic leukaemia cell line THP-1 was maintained in RPMI using the same conditions as described for DMEM. Cryopreserved human hepatocytes from five different donors were obtained from In Vitro Technologies and were maintained in DMEM as described for the cell lines above (paper II). Human primary monocytes were purified from blood obtained from anonymous donors and were cultured in RPMI and 10 % human AB serum or differentiated into macrophages as described in next section (paper IV).

3.1.4 Monocyte isolation and macrophage differentiation (paper IV)

Monocytes were purified from buffy coats obtained from unknown healthy blood donors by endotoxin-free Ficoll purification (Ficoll paque PLUS, GE Healthcare) followed by two steps of Percoll purifications (Percoll PLUS, GE Healthcare) as described in Repnik et al [179], which resulted in about 70 % CD14 positive cells, as determined by FACS analysis. The monocytes were seeded out in 6-well plates in RPMI containing 100 ng/ml M-CSF, 20% human AB-serum, penicillin (100 U/ml) and streptomycin (100 µg/ml). After 7 days, the monocytes had differentiated into macrophages and the medium was replaced with RPMI containing 10 % hAB serum and antibiotics as above (M0). To obtain M1 macrophages, 100 ng/ml LPS and 20 ng/ml IFN γ was also added to the fresh culture medium, whereas for M2, Il-4 was added at a concentration of 20 ng/ml. After 6, 24 and 72 hours, the cells were harvested for RNA using Qiazol or saved for FACS analyses.

For the monocyte studies, the EasySep human monocyte enrichment kit (#19059, Stem Cell Technologies) was used straight after the Ficoll purification. The cells (>90 % CD14 positive cells as determined by FACS analysis) were seeded out at in 6-well plates in RPMI containing 10% human AB-serum, penicillin (100 U/ml) and streptomycin (100 µg/ml), in the presence or absence of 100 ng/ml LPS. The cells were harvested after 8 or 24 hours, using Qiazol (Qiagen) for subsequent analysis of total RNA, including miRNA.

3.1.5 Flow Cytometry Analysis (paper IV)

Differentiated macrophages (M0, M1 and M2) were incubated with APC (allophycocyanin)/Cy7-conjugated antibody CD206 (BioLegend). Analyses were performed using a CyAn ADP Analyzer flow cytometer and the Summit software (Beckman Coulter).

3.1.6 DNA sequencing (paper I-IV)

Sequencing was performed using a Big Dye Terminator Kit and an automatic sequencer (Genetic Analyzer 3100, Applied Biosystems) to control that all cloned constructs were correctly assembled and to verify the mutations introduced after *in vitro* mutagenesis.

3.1.7 Plasmids and expression vectors (paper I-IV)

- **Cloning** of promoters was performed using PCR amplification, followed by subcloning into the PCR 2.1 vector (Invitrogen) and subsequent transfer into the reporter gene vector pGL3basic (Promega) (paper I-III). Likewise, approximately two kb of the PPARD 3'-UTR was cloned into the psiCHECK2 vector (Promega) for subsequent use in the miRNA reporter assays. The 36 bp constructs covering the miRNA target sites of PPARD were obtained by annealing complimentary oligonucleotides for the sites and subsequent cloning into the psiCHECK2 vector (paper IV).

- **In vitro mutagenesis** of putative PPREs or miRNA target sites was performed using oligonucleotides containing the desired mutations and a mutagenesis kit from Stratagene.

- The **expression vector** for human PPARA was a kind gift from Eleanor S Pollak, University of Pennsylvania School of Medicine, PA, USA. The expression vector for PPARD was kindly provided by Dr. CN Palmer and is described elsewhere [180]. The PPARD expression vector was modified into PPARD2 by replacement of the 3'-end of PPARD1 with a PCR-product containing the 3'-end of PPARD2 (paper III). To obtain the full-length PPARD, the whole 3'-UTR of PPARD1 was inserted into the vector coding for PPARD1 (paper IV).

- The **TNT vector** coding for PPARD was cloned as described above, using the full-length sequence for PPARD. The TNT vectors for PPARA, PPARG and RXRA were all cloned by our collaborators at Astra Zeneca.

- The **miRNA expression vectors** coding for either miR-9, miR-29 or miR-155 were created by PCR amplification of the genomic loci and subsequently cloned into the pcDNA 3.1/V5-His-TOPO, by our co-authors at CCK, Karolinska Institutet.

3.1.8 Transient transfection and Luciferase assays (paper I-IV)

For the reporter assays, the cells were seeded out in 24-well plates in PEST-free medium and the following day, the appropriate reporter constructs and expression vectors were added together with lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions. For the promoter studies, agonists or other compounds were added in fresh medium after 24 hours and the cells were treated for an additional 24 hours, whereas the miRNA reporter assays were harvested after 24 hours of transfection. The cells were then lysed with a triton-X containing lysis buffer and the

luciferase activities of the lysates were measured in a luminometer (Lucy2, Anthos). For normalisation of the transfection efficiency, a β -galactosidase plasmid was cotransfected with the reporter plasmid and β -galactosidase activity was also measured (paper III). The psiCHECK2 vector used in paper IV contains the sequences coding for both renilla and firefly luciferase, hence no external normalisation plasmid was needed and a dual luciferase assay, measuring both renilla and firefly luciferase was run for these plasmids. In paper I and II, no normalisation was used for transfection efficiency, however, the transfections were repeated several times with similar results.

3.1.9 RNA isolation, reverse transcription and quantitative real-time PCR (paper I-IV)

Total RNA isolation was performed using the RNeasy system (Qiagen) and reverse transcription was performed on 350-1000 ng of total RNA using a poly-dT primer and Superscript II or III (Invitrogen) according to the manufacturer's instructions. Real-time PCR was performed using gene-specific TaqMan assays (paper I, II, IV) or PPARD exon-specific primers and probes (paper III). The reaction conditions involved an initial denaturation step for 10 min at 95°C, followed by 45 cycles of amplification with 15 sec at 95°C and 1 min at 60°C, using an ABI prism 7000 (Applied Biosystems). All samples were analysed at least in duplicates and the data was analysed using the comparative Ct-method.

For miRNA detection, the miRNeasy kit (Qiagen) was used for isolation of total RNA. Reverse transcription was performed on 10 ng total RNA using specific primers for miR-9, miR-29 or U48 together with the TaqMan miRNA reverse transcription kit (Applied Biosystems). The cDNA was amplified by real-time PCR with the corresponding TaqMan miRNA assays (Applied Biosystems) according to the instructions, using the conditions described above.

3.1.10 Coupled transcription/translation (paper I-III)

Coupled *in vitro* transcription and translation was performed to determine relative protein expression from different mRNA species (paper III) and to obtain the proteins used in the EMSAs (papers I-III). The TNT Quick coupled Transcription/Translation system (Promega) was used together with 1 μ g of the TNT plasmid and either [³⁵S]methionine (Amersham) or unlabelled methionine according to the manufacturer's

instructions. The translation products were subsequently separated by a 10 % SDS-PAGE, subjected to autoradiography and analysed on a PhosphoImager (paper II) or used in the EMSAs as described below.

3.1.11 Western blot (paper II-IV)

Whole cell extracts from primary hepatocytes treated with fenofibric acid (paper II) were subjected to Western blot analysis under reducing conditions as described by Laemmli [181], using an ALT1 specific antibody (paper II). Likewise, nuclear extract from HeLa cells transfected with PPARD1, PPARD2 or empty expression vector (paper III), or from monocytes treated with LPS or vehicle (paper IV), were subjected to Western blot analysis using a PPARD specific antibody, detecting the N-terminal part of the protein (sc-7197, Santa Cruz). Furthermore, *in vitro* translated PPARD from the different 5'-variants in paper III were subjected to Western blot analysis using an antibody detecting the C-terminal of PPARD (IMG 3297, Nordic Biosite). In paper IV, nuclear extracts from primary monocytes treated with LPS or vehicle for 24 hours were subjected to Western blot analysis, using precasted gradient gels 4-12 % (Criterion XT Bis-Tris Gels, BioRad) and the PPARD antibody sc-7197 (Santa Cruz).

3.1.12 Electrophoretic mobility shift (EMSA) (paper I-III)

EMSA was performed to study binding of the PPARs to putative PPRES (paper I and II) and also to study the binding of PPARD1 and PPARD2 to a classical PPRES (paper III). Double-stranded oligonucleotides corresponding to the putative PPRES, or to the classical rat acyl-CoA (ACO), were labelled with [γ -³²P] and mixed with a buffer containing 2 μ g poly(dI-dC), 0.75 mM EDTA pH 8.0, 18 mM HEPES pH 7.9, 0.5 mM dithiothreitol and 4 % Ficoll and *in vitro* produced PPAR and RXR in a molar ratio of 3:1. For competition, 100-fold molar excess of unlabelled double-stranded oligonucleotide was added and for the supershift, the PPARD antibody sc-7197 was used. The mixtures were subjected to a 6 % polyacrylamide gel which was run 4h at 200 V, dried and then analysed on a PhosphoImager.

3.1.13 Chromatin immunoprecipitation (ChIP) (paper I-II)

ChIP was performed to assess *in vivo* binding of PPARD to the promoters of apoA-II and ALT1 (paper I and II). HepG2 cells were treated with PPAR agonists for 8 hours and the DNA and protein were subsequently cross-linked using formaldehyde. The chromatin was purified, followed by enzymatic digestion and immunoprecipitation

using a PPARA, G and D antibody, respectively, or a control IgG. Immunoprecipitated DNA was subjected to PCR with primer pairs specific to the putative PPREs in apoA-II and ALT1, respectively, whereas primer pairs amplifying regions further downstream were used as negative controls. Chromatin input was used as a positive control while H₂O was used as blank control templates for both primer pairs.

3.1.14 Rapid amplification of cDNA ends (RACE) (paper II)

Both 5'- and 3'-RACE were performed to identify possible alternative 5'-ends in transcripts of PPARD and to study alternative 3'-splicing. Marathon cDNA from placenta, adipose tissue and pancreas (Clontech) were used together with the adapter primers AP1 and AP2, together with combinations of PPARD-specific primers. Two sequential rounds of either 5'- or 3'-RACE were carried out and subsequently the obtained PCR products were purified and sequenced. Additional rounds of nested PCR were performed with the AP2 primer and exon-specific PPARD primers in order to enrich for transcripts expressed at low levels.

3.1.15 Human subjects (paper II)

The clinical study was a phase 1 study with the intentions to assess safety, tolerability, effects on lipids and pharmacokinetics of repeated oral doses of the PPARA agonist AZD4619. The study included 20 healthy males randomized to receive once-daily doses of 5 mg AZD4619 (n=15) or placebo (n=5) for 21 days. The ages of the participants were ranged from 20-29 years and their BMI were between 20 to 27.

3.1.16 Statistical Analysis (paper I-IV)

In order to establish differences in reporter vector activity or miRNA/mRNA expression between control and treated cells, Student's t-tests (two-tailed) were used (papers I-IV). For differences in the variables measured over time in the clinical data in paper II, the paired t-test was used. Significant differences are indicated as: *p<0.05, **p<0.01, ***p<0.001.

4 RESULTS AND DISCUSSION

4.1 PPARD activation increases expression of the human apolipoprotein gene (paper I)

ApoA-II is one of the major apolipoproteins of the HDL particle but its anti-atherogenic role is not as established as for apoA-I [10]. However, recent data show that apoA-II is associated with a decreased risk of CVD [182]. The PPARA activating fibrates are shown to increase the levels of both apoA-I and apoA-II in plasma in humans, as well as the HDL cholesterol, supporting the role of apoA-II as anti-atherogenic. Moreover, obese rhesus monkeys treated with the PPARD agonist GW501516 show elevated plasma levels of apoA-I and apoA-II, as well as increased HDL cholesterol levels, further suggesting a beneficial role of apoA-II [95]. It is established that the effect of the fibrates on the apoA-II gene is mediated through a PPRE in the proximal promoter located at -737/-717, denoted the J-site [84]. In contrast, there is no information about the effects of PPARD activation on the apoA-II gene, hence we decided to investigate this matter. Treatment of HepG2 cells with the PPARD agonist GW501516 increased expression of apoA-II mRNA by 1.22-fold as measured by quantitative real-time PCR. Transient transfections of a 3 kb promoter construct of apoA-II in both HepG2 and HuH-7 cells showed that the effect was due to increased transcription and that a cotransfected PPARD plasmid potentiated this effect. The apoA-II promoter was analysed *in silico* to identify putative PPRES and, additional to the J-site, one putative PPRES was found at -2656/-2636, denoted the O-site. This site contained an equal number of identical nucleotides of the core PPRES consensus as the J-site. Furthermore, the O-site contained 4/7 matches in the 5'-flanking region compared to only 2/7 in the J-site (Fig. 11). Interestingly, sequential mutagenesis of these two PPRES in the promoter revealed that the effect of the PPARD agonist was mediated only through the J-site and not the O-site.

	<u>5'-flank</u>	<u>DR1</u>	
Consensus	c ^a a ^a a ^a t ^a ct	AGGTCA ^a AGGTCA	
O-site	caagtggGGGTGAgAGATCA		4/7 + 9/13
	*** * *** * * * *		
J-site	ttctaccAGGGTaaAGGTTG		2/7 + 9/13
	** *** * * * * *		

Figure 11. Comparison of the putative PPRES in the apoA-II promoter

Comparison of identified PPRES in the apoA-II promoter designated as the O-site and J-site, respectively, and the consensus PPRES sequence. The number of matches for the 5'-flanking region and the core DR1, respectively, compared to the consensus sequence are shown to the right. Identical nucleotides between the consensus sequence and the PPRES are denoted by an asterisk.

As previously reported [84] not only PPAR-induced activity was abolished when the J-site was mutated but also the basal activity decreased, reflecting the fact that the J-site is an important regulatory region also for other members of the nuclear receptor superfamily [183]. Both electromobility shift assays (EMSA) and chromatin immunoprecipitation assays (ChIP) confirmed the binding of PPARD to the J-site, establishing the role of apoA-II as a PPARD target gene. Increased transcription of the apoA-II gene is relevant for the abundance of apoA-II in plasma, since the apoA-II levels are controlled mainly by its synthesis rather than catabolism [184]. Whether the effects of an increase in apoA-II levels after PPAR agonist treatment adds on to the effects of the concurrent increase in apoA-I remains to be elucidated. The average apoA-I/apoA-II molecular ratio in human plasma is 2:1, which suggest that such an abundant protein should have important physiological functions [185]. Since increased synthesis of apoA-II is correlated with elevated plasma HDL concentrations, apoA-II might be one of many genes responsible for the beneficial effects regarding lipid and lipoprotein metabolism observed as a result of PPARD activation.

4.2 PPAR agonists regulate the ALT1 gene expression in human hepatocytes (paper II)

Even though ALT activity is used as a standard marker for hepatotoxicity in human clinical trials of pharmaceutical drugs, not much is known about the regulation of the two ALT genes, GPT1 and GPT2. The genes give rise to ALT1 and ALT2, respectively, which have similar enzymatic activities. ALT1 is highly expressed in liver, skeletal muscle and kidney, whereas ALT2 is found in skeletal muscle and heart but not in the liver and kidney [44]. Hence, ALT1 is the dominant isoform expressed in

the liver and it has been shown to be responsible for the basal ALT activity of human normal serum [44]. In a phase I clinical trial of AZD4619, a new PPARA activating compound developed by Astra Zeneca, the serum ALT levels increased over time in some of the participants. The trial went on for 21 days and at the last day of dose, five of the 15 participants had ALT levels above upper level of normal (ULN), defined as the 95 percentile of ALT activities from pre-dose (Fig. 12). At follow-up at day 31, seven of the participants were above ULN. Similarly, AST-levels increased in some of the participants during the study but no other liver markers, such as γ -glutamyl transferase, t- bilirubin, alkaline phosphatase, pro-thrombin or creatine kinase, were elevated. However, due to the increased ALT and AST levels, AZD4619 was discontinued from development.




Figure 12. Serum ALT activities in healthy volunteers treated with AZD4619

15 human subjects were treated daily with 5 mg AZD4619 for 21 days. Serum ALT activities (U/l) increased after treatment with AZD4619 in some of the 15 subjects. ULN = upper level of normal

Since previous *in vitro* studies have shown that the AST gene is transcriptionally regulated by PPARA [186] we focused on testing the hypothesis that PPAR agonists regulate ALT gene expression. The PPARA agonist fenofibric acid was used as a model compound for PPARA activation and increased ALT1 expression in human primary hepatocytes compared to untreated cells, as measured by Western blot. Fenofibric acid treatment also increased mRNA expression of ALT1 in three out of five different donors of primary hepatocytes measured by quantitative real-time PCR, indicating that the effect varies between individuals. Treatment of the hepatoma cell

line HuH-7 with fenofibric acid showed that ALT1 but not ALT2 mRNA expression was induced after the treatment compared to untreated cells. To establish if the effect was transcriptional, two kb of the GPT1 and GPT2 promoters, respectively, were cloned into a reporter vector and subsequently used in transient transfection assays in HuH-7 cells. Both fenofibric acid and AZD4619 showed dose-dependent increases of ALT1 promoter activity, whereas the ALT2 promoter did not respond. Thus, the effects of PPARA on ALT enzyme activity are accounted for by ALT1. Furthermore, PPARG and PPARD agonists were also shown to activate the ALT1 promoter, indicating that all the PPARs had similar effect on the promoter. In contrast, other potent compounds such as 12-O-Tetradecanoylphorbol 13-acetate (TPA) and TNF α did not activate the ALT promoter (Fig 13).




Figure 13. Induction of ALT1 promoter by PPAR agonists in transfection assays

PPARA agonists: Fenofibric acid (FA), Clofibrate (clofib.); PPARG agonist: Rosiglitazone (Rg); PPARD agonist: GW501516; Miscellaneous: 12-O-Tetradecanoylphorbol 13-acetate (TPA), tumour necrosis factor alpha (TNF- α).

Using *in silico* tools, the promoter of GPT1 was screened for PPRES and seven putative sites were found. EMSA studies verified that PPARG could bind to a site located at -574 in the promoter, whereas it could not bind to the other putative PPRES. Furthermore, mutation of the -574 site abolished the binding of PPARG in the EMSA and reduced the responsiveness to PPAR agonists of the promoter construct. Noticeably, PPARA did not bind to this site using EMSA, even though it could bind to the control rat ACO-PPRE. Importantly, PPARG is known to bind more strongly than the other PPARs to PPRES and the lack of binding of PPARA in this study might be due to suboptimal conditions of the EMSA or to the fact that the PPRE was weak [76].

Since EMSA only reflects binding to naked DNA, a ChIP was performed to evaluate the effects of the PPAR agonists in the presence of chromatin. The ChIP confirmed binding of both PPARA and PPARG to the -574 PPRE. This study shows that PPAR agonists can induce ALT1 by increased transcription of the ALT1 gene which is an alternative explanation for increased ALT protein in plasma.

Since ALT was introduced in clinical monitoring during the 1950s [43], much attention has been given to its role as a hepatotoxicity marker. In contrast, the function of ALT as a metabolic enzyme important for gluconeogenesis has not been given much notice. It is known that plasma ALT and AST levels increase in people experiencing a rapid weight loss, peaking after 2 weeks [187]. The extreme gluconeogenic condition is a possible explanation for the induction of the aminotransferases in the liver during fasting [188, 189]. Elevations in aminotransferase levels are also associated with obesity and the metabolic syndrome [190, 191], even though it is unknown whether it is the obesity per se which affect ALT activity or not. Furthermore, increased plasma ALT levels are shown to predict the incidence of type-2 diabetes and cardiovascular disease [191, 192]. Gluconeogenesis is increased in subjects with type-2 diabetes and the elevated ALT levels might be due to an enzyme induction as the need for aminotransferases in the liver increases [193]. Thus, ALT elevations in plasma might not only reflect a hepatic injury but the metabolic status of the liver might also play a role for plasma ALT levels. In further support of this alternative hypothesis for ALT elevations is the fact that PPARA is important for gluconeogenesis. PPARA knock-out mice show impaired gluconeogenesis as a response to fasting compared to PPARA wild-type mice [82]. Hence, it is likely that activation of PPARA agonists would induce ALT expression in the liver. Furthermore, moderate but transient elevations in ALT and AST levels are seen in approximately 10 % of the patients treated with fenofibric acid without any other signs of liver injury [50, 51]. This individual variation was also observed in our study where only some individuals responding to AZD4619 treatment with increased aminotransferase levels. Also, there was a difference in the response in ALT expression after treatment with fenofibric acid between the different hepatocyte donors. The PPARG agonists have never been associated with transaminase elevations in the clinic, unless they have been proven to be toxic, as in the case with Troglitazone. Instead PPARG agonists are shown to reduce ALT activity in serum [194], probably due to the fact that they improve the insulin sensitivity, which reduces gluconeogenesis in the liver [195]. The alternative hypothesis of ALT induction by drugs influencing

metabolism was recently demonstrated to be valid also *in vivo* [196]. Rats treated with dexamethasone for 24 hours showed increased levels of ALT activity in the serum and in the livers compared to untreated rats, without any signs of tissue damage within this time. The dexamethasone treated rats also showed increased amounts of glycogen in their livers, indicating an increased gluconeogenesis, which reflects the condition induced by the treatment.

Of note, the moderate and transient increase in ALT activity induced by the PPAR agonists should be distinguished from the massive increases that occur during liver damage. The development of additional hepatotoxicity markers would be important for use in combination with ALT so that the new PPAR compounds have a chance to pass the clinical monitoring of today.

4.3 Regulation of PPARD by alternative splicing (paper III)

Since the majority of studies on PPARD concern the effects of its activation, we decided to investigate its regulation and more specifically; whether it is subjected to alternative splicing, which might give rise to additional isoforms and/or influence the expression. The human PPARD gene consists of 9 exons of which exons 1-3, the 5'-end of exon 4 and the 3'-end of exon 9 are untranslated [92]. In this study, five new alternative exons were identified in the PPARD gene by 5'-RACE using cDNAs from placenta, adipose tissue and pancreas and these exons make up a variable range of PPARD mRNA species (Fig.14). TaqMan analysis using a panel of human cell lines and tissues revealed that the most common splice variant encompassed exon 2 connected to exon 4 (Fig 14, B), whereas the other splice variants were expressed at much lower levels. *In vitro* transcription/translation and Western blot analysis showed that the various transcripts were translated into PPARD protein with different efficiencies, inversely correlated to the length, as well as the number of AUGs in the 5'-UTR. Analysis of basal activity of the previously identified promoter upstream of exon 1 and the alternative promoters upstream of the newly identified putative transcription start sites (Fig 14, H-K) in transient transfection assays, revealed that the promoter upstream of exon 1 is the major region for transcriptional activation.




Figure 14. A schematic representation of the PPARD gene

Coding exons (*white* boxes), previously reported untranslated exons or part of exons (*grey* boxes), and herein identified untranslated exons (*black* boxes). The variety of splicing among untranslated exons and alternative 5'-ends identified by 5' RACE is shown below the gene (5'-UTRs: A-K).

Moreover, a 3' splice transcript encoding a truncated isoform of PPARD, PPARD2, was identified in placenta and adipose tissue. This truncated variant is formed due to intron retention of intron 8 which introduces an early stop codon, leading to a truncated PPARD protein lacking 82 amino acids of the C-terminal containing the ligand-binding domain. Analysis of PPARD2 in transient transfection assays revealed that it could not be transactivated by ligand-binding but rather repressed ligand-induced activation of PPARD1, constituting a dominant negative inhibitor of the full-length receptor. Furthermore, PPARD2 was unable to bind to a classical PPRE in EMSA studies, suggesting that it does not compete with PPARD1 for DNA-binding as is suggested for the dominant negative full-length mouse orthologous of PPARD (E411P) [197, 198]. The mechanism of inhibition of PPARD2 might instead be due to competition for cofactors or its heterodimerization partner RXR, which has been speculated as the negative regulatory mechanism for some of the other truncated nuclear receptors [132, 133, 135, 136].

The diversity in PPARD 5'- transcripts described in this study adds another level of regulation. Post-transcriptional regulation by 5'-splicing have already been described for the mouse PPARD [199] and also for the human PPARG and PPARG genes [200, 201] where the PPARG transcripts also have shown to influence translational efficiency [202]. The mechanism which accounts for differences in translational efficiency between the 5'-splice variants of PPARD is not clear but the fact that long 5'-UTRs and number of AUGs play a role indicate that secondary structures of the 5'-UTR or upstream open reading frames (uORF) might be involved [203]. A study regarding

alternative 5'-transcripts of PPARG identified that the stability of the 5'-UTRs is inversely related to the translational efficiency [202]. Even though alternative promoters have been described to be functional in the mouse PPARG gene [199], the promoter analysis in this study showed low activity compared to the one already described. However, it is possible that the promoters described here might be functional in tissues other than the ones tested in this study, due to differential expression of cofactors. Truncated isoforms have been described for the other members of the PPAR family and for many of the other nuclear receptors. The mechanism of inhibition of PPARG2 remains to be elucidated but general mechanisms suggested for the other nuclear receptors are; competition for binding to the DNA, competition for cofactors/auxiliary factors or for heterodimerization partners, or influences on nuclear localisation [132-136]. The conclusion of this study is that alternative splicing of human PPARG could constitute an intrinsic role for the regulation of PPARG expression and activity, which could be of relevance both in physiology and disease.

4.4 Regulation of PPARG by miRNA (paper IV)

In order to further investigate the mechanisms governing the regulation of the human PPARG gene, the 3'-UTR of PPARG was analysed *in silico* for miRNA binding sites using the UCSC genome browser. Two putative miRNA target sites were identified, miR-9 and miR-29 (Fig 15).




Figure 15. Schematic representation of the PPARG 3'-UTR luciferase reporter construct
Putative target sites for miRNAs and their corresponding miRNA sequences are shown below.

Transient transfection assays in HEK293 cells using a PPARG 3'-UTR reporter construct showed decreased luciferase expression when miR-9 was cotransfected, whereas miR-29 or the control miR-155 did not alter the luciferase activity, indicating that miR-9 can regulate the PPARG gene. Furthermore, when the miR-9 target site was mutated, the effect of miR-9 was abolished, demonstrating that the regulation of the 3'-UTR by miR-9 was mediated through the *in silico* identified miR-9 site (Fig 16).




Figure 16. The PPARD gene is a direct target of miR-9

Luciferase activity of reporter constructs that contain the wild-type (grey bars) of the entire PPARD 3'-UTR or mutated at the putative miR-9 binding site (black bars). HEK293 cells were transiently cotransfected with the PPARD 3'-UTR reporter constructs and expression vectors coding for miR-155, miR-9 or miR-29 as indicated.

Moreover, a luciferase reporter containing only the short sequence coding for the miR-9 target site (36 bp) also showed decreased luciferase activity after coexpression of miR-9, whereas mutation of the seed sequence abolished the luciferase activity. Also, cotransfection of an antisense of miR-9, antagomiR-9, in the absence of any cotransfected miR-9, showed an increase in luciferase activity in cells transfected with the PPARD 3'-UTR or the short miR-9 construct. Furthermore, when miR-9 was transfected into the cells, the luciferase activity could be restored to almost normal levels by antagomiR-9. These results strongly support the finding that miR-9 represses PPARD protein expression. In contrast, miR-29 did not affect the expression of the full-length 3'-UTR PPARD reporter, indicating that it is not a regulator of PPARD. To elucidate whether miR-9 targets only the PPARD protein levels or if mRNA levels also are affected, miR-9, miR-29 or the empty plasmid was transfected into HEK293 cells for 24 h and PPARD mRNA expression was measured by quantitative real-time PCR. However, no difference between the groups could be detected which suggests that miR-9 does not affect the levels of the PPARD transcript, instead it inhibits the translation of the mRNA into protein.

Since miR-9 has been shown to play an important role in the inflammatory response in monocytes [204], the relevance of miR-9 regulation in relation to PPARD expression was investigated in monocytes and macrophages. Based on the knowledge that treatment of primary monocytes with LPS increases miR-9 expression [204], human primary monocytes were treated with LPS to study the effects on PPARD function. Accordingly miR-9 expression as well as PPARD mRNA expression was increased

after both 8 and 24 hours treatment. However, the PPARD target genes adipose differentiation-related protein (ADFP), adipose triglyceride lipase (ATGL) and carnitine palmitoyltransferase (CPT1) were notably lower after 8 hours treatment and only increased after 24 hours, suggesting that the increased expression of miR-9 in the cells was counter-acting the effects of the PPARD elevation. Western blot analysis of monocytes treated with LPS for 24 hours did not show any major differences after LPS treatment compared with cells only treated with vehicle, further proposing that PPARD protein levels were not as increased by the LPS treatment as the PPARD mRNA expression.

Due to the recent identification of PPARD as a major regulator of the M2 phenotype of macrophages, the involvement of miR-9 in the alternative activation of macrophages was elucidated. Human primary monocytes were differentiated into macrophages and subsequently treated with LPS and IFN γ to become M1 macrophages, IL-4 to become M2 macrophages or vehicle to serve as control (M0) for 6, 24 or 72 h. Of note, PPARD mRNA expression was higher in M1 compared to M2 macrophages at 24 and 72 hours, whereas mRNA expression of PPARG was higher in M2 compared to M1 at all time points measured. Analysis of miR-9 expression revealed no statistical difference between the treatments suggesting that miR-9 expression is not involved in the development of different macrophage phenotypes.

MiR-9 is a microRNA which has been reported to be involved in distinct processes such as neuronal differentiation in the brain, release of insulin from the pancreas and the immune response in monocytes [204-206]. By increasing concomitantly with NF κ B in monocytes and targeting its 3'-UTR, miR-9 inhibits the inflammatory cascade and constitutes a negative feed-back loop of the inflammatory response [204]. Similarly, this study shows that PPARD gene expression is increased in monocytes during inflammatory stimulation with LPS and the protein expression is partly inhibited by the simultaneous increase in miR-9. The importance of PPARD for the inflammatory status of macrophages is not clear but recent studies in murine macrophages suggest a function of PPARD in inflammation, by causing a switch from M1 to M2 phenotype [29, 30]. However, in this study PPARD expression was higher in M1 than M2 macrophages, which supports data from a recent human study [31], highlighting the differences between human and mouse macrophages. The lack of differences in

miRNA expression between the macrophage subtypes suggests that the function of miR-9 is of importance only in monocytes but not in differentiated macrophages.

5 GENERAL DISCUSSION

PPARD is a protein with multiple functions in basically all the tissues of the human body. When the PPAR family was discovered in the early 1990s, PPARD was not considered to be more than a house-keeping gene in the shade of PPARG and PPARG, the important mediators of the fibrates and glitazones, respectively. Today, when detrimental side-effects associated with some of the PPARG agonists have been shown and the role of PPARD as a nuclear receptor with beneficial effects on both lipid metabolism and insulin sensitivity have become more clear, a number of studies regarding its functions have emerged. The increased oxidative capacity in muscle, increased fatty acid combustion in adipose tissue and beneficial effects on lipoprotein metabolism are clear indicators of overall positive effects mediated by PPARD, which could be taken advantage of in the treatment of obesity and type-2 diabetes. However, caution has to be taken since PPARD has been reported to increase cell proliferation and to be involved in cancer formation in the colon, even though the literature on the subject is not totally conclusive [207] .

The first two studies of this thesis further supports the role of PPARD as an important regulator of human lipoprotein composition and overall metabolism. The increased expression of apoA-II mediated by transactivation of PPARD is likely one of the reasons for the increase in the number of HDL particles and the enhanced reverse cholesterol transport in primates and humans treated with GW501516. The activation of the ALT1 gene in the liver by all three PPARs implies that increases in lipid oxidation results in a state of the body that favors gluconeogenesis including the enzymes implicated in that process. Hence, PPAR activation mimics a state of starvation of the body which could be taken advantage of in obesity and obesity-related disorders such as type-2 diabetes and cardiovascular disease. However, if the ALT enzymes are to be assessed during clinical trials of developing PPAR agonists, an awareness of the properties of the PPARs to induce aminotransferases by the non-toxic mechanism is important.

The effects of PPARD have mainly been investigated using activation of the receptor by a specific ligand, or by studying the effects of PPARD knock-out or transgenic mice models. These studies show the importance of PPARD activation in lipid metabolism

but have also identified an inflammatory role of the receptor in macrophages. As demonstrated by the LDL receptor knock-out mouse model, a reduction in plaque size is seen both after knock-out of myeloid PPARD and after PPARD activation, due to its sequestering of inflammatory corepressors in the unliganded state [73]. Hence, the effects of regulation of the gene are complex, since not only abundance of the protein but also availability of ligands is important for its downstream effects. Additionally, the availability of coactivators and corepressors, as well as availability of the other PPAR family members or other nuclear receptors might determine the response of PPARD expression or activation [71, 72]. Hence, studies of this nuclear receptor have to be performed in a well controlled experimental setting. No detailed investigations on the regulation of the human PPARD gene has been performed until now, even though the mouse gene has been reported to be regulated by alternative promoter usage and alternative splicing [199]. In this thesis, human PPARD is shown to be subjected to alternative splicing in the 5'-end, which results in differences in translational efficiency between the different transcripts. Furthermore, a 3'-truncated PPARD isoform, PPARD2, is shown to be a negative regulator of the full length variant. These findings could be of importance for PPARD expression in different tissues or during different metabolic conditions, however, the functional relevance of the splice variants remains to be identified. Furthermore, the mechanism of negative regulation of PPARD2 still remains to be elucidated.

Regulation of PPARD by miRNA was also identified as a posttranscriptional regulatory mechanism of the human PPARD gene. However, to understand the importance of PPARD regulation by miR-9, further studies of PPARD function in monocytes is needed. Since miR-9 is highly expressed in the brain and PPARD expression is important for myelination of the corpus callosum [100], the brain might be one of the main sites of PPARD regulation by miR-9. Additionally, the concomitant increase of PPARD and miR-9 in monocytes after inflammatory stimulation suggests a tight relationship between these two players during the initial inflammatory response of monocytes, and might be relevant in the initiation of the atherosclerotic process.

The future for PPARD as a drug target is not easy to predict. With the declining interest for the PPARA and PPARG agonists on the market, a PPARD compound might take over as their replacement therapy. However, the aminotransferase elevations of PPARA agonists in clinical trials as well as the failures of some of the PPARG agonists which

have been on the market for many years, the PPARs in general have a bad reputation. Also, non-resolved issues about its involvement in tumour formation are of major concern, even though there has been no indication of increased tumourogenesis in the millions of patients taking fibrates or glitazones. Another obstacle for the development of PPAR agonists is the fact that all new PPAR ligands require two years of preclinical oncogenicity studies prior to initiating any clinical studies of 6 months and longer, which might not make the effort worth-while for pharmaceutical companies [118]. However, tissue-specific PPAR modulators (SPPARMS) might be the resolution to this problem. The future might hold a muscle-specific PPARD modulator that will be prescribed additionally to exercise to patients with the metabolic syndrome. The exercise itself would increase PPARD expression in the skeletal muscle and the drug would cause its activation, resulting in increased expression of genes involved in lipid oxidation. The increased oxidation of fatty acids in the muscle increases the overall insulin sensitivity, and is beneficial for the endurance capacity of the muscle fibers. Of note, not the PPARD agonist itself but the combination of exercise and PPARD agonist is important for the beneficial effects, as recently demonstrated by a study of sedentary mice administrated either with GW501516 alone or in combination with an exercise program [208]. Exercise not only increases PPARD expression in skeletal muscle but also improves the metabolic status of the patients, which might be relevant for the synergic function of PPARD activation and exercise. An alternative might be to influence the splicing machinery to only create short 5'-ends of PPARD mRNA, hence increasing the translational efficiency. Whether that is applicable or not is too early to say but the idea is intriguing. Treatment with an antagomiR-9 might also be a way to increase PPARD expression. However, miR-9 have additional targets, both known and unknown which probably would cause numerous side-effects. To be able to modulate this receptor in a clinical setting, it is important to further elucidate by which mechanisms it is regulated. With increasing knowledge about different functions of PPARD, additional studies on the regulation of this nuclear receptor will probably emerge to resolve new treatments for metabolic syndrome and CVD. Until then, exercise in combination with a high dietary intake of PUFAs will be the most effective way to increase the expression of PPARD and subsequently cause its activation, not only to treat but also to prevent the development of the metabolic syndrome.

6 CONCLUSIONS

The work of this thesis has identified new target genes of PPARD and also revealed post-transcriptional mechanisms of regulation of this nuclear receptor. In summary, the conclusions of this thesis are;

- **Paper I:** PPARD activation increases the expression of the human apoA-II gene, which might be one of the reasons why treatment with PPARD agonist increases production of HDL particles and stimulates reverse cholesterol transport.
- **Paper II:** The ALT1 gene is a transcriptional target of the PPARs, which may result in mild ALT elevations in plasma in the absence of liver injury when assessing new PPAR agonists in clinical trials.
- **Paper III:** PPARD is subjected to alternative splicing, both in the 5'- and 3'-ends, which identifies possible mechanisms for PPARD regulation in time and space.
- **Paper IV:** The human PPARD gene is regulated by miRNA-9, which might be important for the early inflammatory response in monocytes.

7 ACKNOWLEDGEMENTS

This research program was conducted at the Atherosclerosis Research Unit, King Gustaf V Research Institute and the Center for Molecular Medicine, Karolinska Hospital during 2005-2011 and was partly supported by the Swedish Heart-Lung Foundation.

During these years I have experienced feelings of disappointment, stress and despair but also excitement and fulfillment. I have been fortunate to have many people supporting me and I would especially like to express my sincere gratitude to:

Ewa Ehrenborg, my main supervisor. Thank you so much for taking me into your group and for believing in me. I really appreciate your enthusiasm and positive spirit, as well as the freedom you have given me during these years. Thank you for always being patient with me no matter what; when I made mistakes in the lab, when I had problems interpreting my results or when you had to correct my writing over and over again. I am also very thankful for your support and understanding when life has been tough.

Björn Glinghammar, my co-supervisor. You have been great support from day one and I feel that you contributed a lot towards my decision to start a PhD. You were an excellent teacher during my first time at GV and you are still just an e-mail away. I really appreciate you sharing ideas on how to become a millionaire, about nutrition and general things about life. I also want to thank you for giving me the opportunity to work on the ALT-project together with you.

Kerstin Lundell, my co-supervisor. Thank you for always helping me out in the lab and for making me realize that organization of tubes and boxes is important. It was always fun working with you and I was sad when you left the group. I appreciate your friendship and I am very grateful that you wanted us to work together on the PPARD paper and share the authorship.

Ian Cotgreave, my mentor. Thank you for accepting to be my mentor with such short notice and for making me realize that writing the thesis should be fun. I hope we can continue the mentoring even after I get my PhD degree.

Anna Aminoff, my PhD companion and lab sister. Thanks for being my close lab partner these years and for being as addicted to good coffee as myself. Thanks also for the company on all the trips during the years and for being such a relaxed individual. I have missed you during your maternity leave. Thanks for also helping me with the PPARD genotyping and statistics.

Olivera Werngren, my fantastic lab technician. Thanks for all the fun times in the cell room - experiencing both triumph and hopelessness together. You have been such great support since you came into the group and the last manuscript wouldn't have been much without you. I am amazed by your abilities to perform so much in a limited time, for taking the results personally and for never giving up. And also for bringing a lunch box every day.

Joelle Magne, our newest group member. Thanks for bringing a Guadeloupian/French touch into the group and for introducing us to delicious macaroons. Also, for sharing thoughts and feelings and for spending evenings at work.

My colleagues in the Cardiovascular Genetics group:

Anders Hamsten, for elegantly leading the big Cardiovascular Genetics group and for always chipping in with your Euros; **Rachel Fisher**, for teaching us about the English style of life: afternoon tea with scones and cucumber sandwiches and traditional fruit cake for Christmas; **Ferdinand van't Hooft**, for your critical questions during our group meetings which have made me extra motivated to study before my Wednesday meeting presentations; **Per Eriksson**, for being such a genuine person and for answering all the lab trouble questions I couldn't understand; **Angela Silveira**, for interesting stories about Brazil and Argentina during wintertime; **Louisa Cheung**, for helping me with the FACS at any time and for teaching me that dried mango is delicious; **Joanna Chmielewska**, for being funny and talkative (in all languages it seems); **Maria Iglesias**, for being "mi vecina favorita", for your friendliness and all our talks about motherhood and life, and for patiently answering all my questions about THP1-cells and LPS; **Sanela Kurtovic**, for bringing me (us) Tim-Tams from your trip to Australia; **Ekaterina Chernogubova**, for sharing thoughts and ideas, and for taking a big responsibility in the cell lab; **Hovsep Mahdessian**, for always asking questions which make me realize that I should know the answers; **Shohreh Maleki**, for introducing me to Finn Crisp and roasted sun seeds; **Valentina Paloschi**, for your charming personality, for inviting me to dinners and for our runs; **Hanna Björck**, for contributing to a nice work atmosphere; **Emina Vorkapic** for being a high-quality student; **Therese Olsson**, for help in the lab and all the friendly chats; **Karin Husman** for ordering tubes and plates on my demand and for your curiosity; **Fariba Foroogh**, for being so sweet and helpful at all times, I really appreciate it!; **Peri Noori**, for teaching me how to use the bioanalyzer in the early days at GV and for coming to our group with your miRNA expertise; **Birgitta Söderholm**, for being an early bird; **Ami Björkholm**, for always being helpful, chatty and friendly; **Alexandra Bäcklund**, for your thoughtfulness and your beautiful accent; **Rona Strawbridge**, for funny stories during lunch time; **Bengt Sennblad**, for being a cross-word expert; **Dick Wågsater**, for keeping the standards of the cake list high; **Karin Danell-Toverud** for always ordering things as fast as possible every time - it has been a great help!; **Karolina Anner**, for being such a sweetheart; **Karl Gertow**, for always being a funny guy and for finding us a funny girl; **Maria Gonzalez Diez**, for your love of the Swedish pop group "the Ark" ☺; **Maria Sabater**, for being such a happy sparkle; **Sarah-Jayne Reilly**, for being such a funny character and running inspiration, and for the help with the English; **Olga Piksasova**, for friendly talks and laughs; **Lasse Folkersen**, for cheering up the atmosphere and for talking to me in English; **Malin Larsson**, for sharing thoughts and ideas, mainly about babies but also about science; **Jacob Odeberg**, for being an expert on gene regulation; **Magnus Mossfeldt**, for helping me transfer my 16 000 e-mails into the new system and for chatting about Agde; **John Öhrvik**, for making statistics more fun.

My colleagues in the Membrane Signaling Networks group:

Alejandro Bertorello, for always smiling to me in the corridor; **Kristina Eneling**, for being a good friend with a warm heart, a chatting mouth and an early “let’s go for lunch-stomach”; **Laura Brion** for contributing to a nice atmosphere in the lab; **Sergej Popov**, for keeping the isotope lab nice and tidy ☺.

Former colleagues:

Josefin Skogsberg for being a great role-model, for nice lunches, for supporting me during the writing of this thesis and for running around Haga together; **Per Sjögren**, for the time at GV, for having an even higher consumption of coffee than me and for challenging me during our runs around Haga; **Karin Stenström**, for your friendship and crazy personality. For always being a step ahead of me in helping to sort practicalities - no matter if it has to do with pregnancy or writing a thesis; **Maria Mannila**, for the time when we were roommates at GV and for making me laugh every time I hear the Abba song: “When I clean the kitchen”. Thanks also for helping me with the PPARD statistics; **Anita Larsson**, for being such a sweet person, for all the help with the ELISAs and for also teaching me how to run them myself; **Barbro Burt**, for all the help in the lab during the years. You are missed by everyone but of course you know that; **Zongpei Chen**, for your excellent guiding through Shanghai and Hangzhou; **Maria Kolak**, for being funny and generous; **Ann Samnegård**, for interesting details on how to perform a PTCA; **Helena Ledmyr**, for help in the lab in the early days; **Katja Kannisto**, for having a really “Porsche” dissertation party; **Massimiliano Ria**, for your Italian style; **Sergey Kraipner**, for your kindness; **Alexander Kovacs** for your funny personality; **Mattias Sjöström** for teaching me that the pinky is vulnerable, no matter how strong you are; **Justo Sierra Johnson** for always being considerate and generous; **Sara Hägg** for your singing skills; **Katrin Brandt** for nice lunches at work and time spent together during our maternity leave; **Therese Edholm** for giving me a ride to Sättra once a week during the GV time; **Camilla Berg** and **Karin Björklund-Jonsson**, for all the nice lunches outside GV in summertime; **Fei Chen**, **Marie Björnstedt-Bennermo**, **Hanna Björk**, **Vincent Fontaine**, **Monzur Kazi**, **Anders Mälarstig**, **Chaoyong Zhu**, **Jacob Lagerkrantz**, **Johan Björkegren**, **Jesper Tegnér**, for making GV a great work place.

My collaborators:

Ingalill Rafter for your help with cloning and EMSAs and for time spent in the isotope lab; **Cecilia Alexandersson**, for fantastic cloning and maxiprep company; **Dan Grandér**, for making time for the miR-9 project and for serving strong coffee; **Martin Corcoran**, for coming up with the idea for the miR-9 project and for allowing me to join you in the lab. You are a true miRNA guru!

Thanks to the people in the Ståhle group for friendly discussions on miRNA related topics and everyday life matters, especially **Andor Pivarcsi**, **Ning Xu**, **Tianling Wei**, **Florian Meisgen** and **Clara Ibel Chamorro**. I also want to thank the members of the Hansson group for your help regarding monocyte differentiation, especially **Lotta Tammik**, **KC**, **Daniel Johansson** and **Hanna Agardh**.

Thanks to everybody at **CMM floor 2** for contributing to a nice work atmosphere.

Thanks to the staff at “**Glada restaurangen**” for the friendly faces at lunch time and excellent pizzas.

Thanks to all my coaches in Hammarby Friidrott, **Petra**, **Magnus**, **Peter** and **Micke G** for helping me during these years even though I was no star.

All my other friends:

Maja, for your true friendship and all our trips together; **Ame**, for always being a good friend even though we don't see each other that often (one story has kept us apart); **Linn** and **Ulrika**, for caring for me and for all our talks, walks, training sessions, cozy "fikas" and dinners; **Lisa**, for everything we got up to in the old days - I hope for more of them; **Robert**, for all our travels and for being a friend; **Jenny**, for all the time spent together during the two last years and for being such a sweetheart; **Berit-Elin** and **Lene**, for all the fun times in Australia and after; **Mathias**, for all the hours of talking in Sättrahallen and for great dinners at Bredängsgrillen; **Johanna** and **Anders**, for making sure my daughter learns Norwegian traditions and for babysitting during my dissertation party; Årstagänget: **Peter**, **Hampus**, **Irina**, **Jens W** and **Jens D**, we hardly see each other anymore but I do enjoy the occasions when we manage to get together. I hope you can all come to my party!

My family:

To my parents; **Paula** and **Rickard**, for all the support in my everyday life and for going away during my thesis writing so that I could have my own mansion; my sister **Marie**, for all the babysitting and time spent together; my brother **Micke**, for making me feel like a big sister; auntie **Lotta**, in Australia for being my extra mum during all my time spent there.

To **Chris**, for being the love of my life and for all the support during the writing of this thesis and to my daughter **Lea**, for making me realize that nothing in life could be more important. I love you both so much ♥

Petra

Stockholm, January 23, 2011



"What fits your busy schedule better, exercising one hour a day or being dead 24 hours a day?"

(Adapted from www.hapi-naples.com)

8 REFERENCES

1. Mathers, C.D., T. Boerma, and D. Ma Fat, *Global and regional causes of death*. Br Med Bull, 2009. **92**: p. 7-32.
2. Luchsinger, J.A., *A work in progress: the metabolic syndrome*. Sci Aging Knowledge Environ, 2006. **2006**(10): p. pe19.
3. Zimmet, P., K.G. Alberti, and J. Shaw, *Global and societal implications of the diabetes epidemic*. Nature, 2001. **414**(6865): p. 782-7.
4. WHO, <http://apps.who.int/bmi/index.jsp>.
5. van Dieren, S., et al., *The global burden of diabetes and its complications: an emerging pandemic*. Eur J Cardiovasc Prev Rehabil, 2010. **17 Suppl 1**: p. S3-8.
6. Enkhmaa, B., et al., *Postprandial lipoproteins and cardiovascular disease risk in diabetes mellitus*. Curr Diab Rep, 2010. **10**(1): p. 61-9.
7. Champe, H., *Lippincott's Illustrated Reviews: Biochemistry 2nd edition*. 2nd ed. 1994.
8. Smirnov, A.N., *Lipid signaling in the atherogenesis context*. Biochemistry (Mosc), 2010. **75**(7): p. 793-810.
9. Forti, N. and J. Diament, *High-density lipoproteins: metabolic, clinical, epidemiological and therapeutic intervention aspects. An update for clinicians*. Arq Bras Cardiol, 2006. **87**(5): p. 671-9.
10. Tailleux, A., et al., *Apolipoprotein A-II, HDL metabolism and atherosclerosis*. Atherosclerosis, 2002. **164**(1): p. 1-13.
11. www.invivo.ca/illustration_and_print_formation_of_atherosclerosis.html. Copyright Resverlogix Corp.
12. Hansson, G.K. and P. Libby, *The immune response in atherosclerosis: a double-edged sword*. Nat Rev Immunol, 2006. **6**(7): p. 508-19.
13. Ross, R., *Atherosclerosis--an inflammatory disease*. N Engl J Med, 1999. **340**(2): p. 115-26.
14. Napoli, C., et al., *Fatty streak formation occurs in human fetal aortas and is greatly enhanced by maternal hypercholesterolemia. Intimal accumulation of low density lipoprotein and its oxidation precede monocyte recruitment into early atherosclerotic lesions*. J Clin Invest, 1997. **100**(11): p. 2680-90.
15. Chan, D.C. and G.F. Watts, *Dyslipidaemia in the metabolic syndrome and type 2 diabetes: pathogenesis, priorities, pharmacotherapies*. Expert Opin Pharmacother, 2011. **12**(1): p. 13-30.
16. Berneis, K.K. and R.M. Krauss, *Metabolic origins and clinical significance of LDL heterogeneity*. J Lipid Res, 2002. **43**(9): p. 1363-79.
17. Cybulsky, M.I. and M.A. Gimbrone, Jr., *Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis*. Science, 1991. **251**(4995): p. 788-91.
18. Nakashima, Y., et al., *Upregulation of VCAM-1 and ICAM-1 at atherosclerosis-prone sites on the endothelium in the ApoE-deficient mouse*. Arterioscler Thromb Vasc Biol, 1998. **18**(5): p. 842-51.
19. Rajavashisth, T.B., et al., *Induction of endothelial cell expression of granulocyte and macrophage colony-stimulating factors by modified low-density lipoproteins*. Nature, 1990. **344**(6263): p. 254-7.
20. Ottend, E., et al., *A macrophage receptor for oxidized low density lipoprotein distinct from the receptor for acetyl low density lipoprotein: partial purification and role in recognition of oxidatively damaged cells*. Proc Natl Acad Sci U S A, 1995. **92**(5): p. 1391-5.
21. Hansson, G.K., et al., *T lymphocytes inhibit the vascular response to injury*. Proc Natl Acad Sci U S A, 1991. **88**(23): p. 10530-4.
22. Newby, A.C., *Metalloproteinase expression in monocytes and macrophages and its relationship to atherosclerotic plaque instability*. Arterioscler Thromb Vasc Biol, 2008. **28**(12): p. 2108-14.
23. Bennett, M.R., *Reactive oxygen species and death: oxidative DNA damage in atherosclerosis*. Circ Res, 2001. **88**(7): p. 648-50.

24. Branen, L., et al., *Inhibition of tumor necrosis factor-alpha reduces atherosclerosis in apolipoprotein E knockout mice*. Arterioscler Thromb Vasc Biol, 2004. **24**(11): p. 2137-42.
25. Shimada, K., *Immune system and atherosclerotic disease: heterogeneity of leukocyte subsets participating in the pathogenesis of atherosclerosis*. Circ J, 2009. **73**(6): p. 994-1001.
26. Bouhrel, M.A., et al., *PPARgamma activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties*. Cell Metab, 2007. **6**(2): p. 137-43.
27. Stout, R.D., et al., *Macrophages sequentially change their functional phenotype in response to changes in microenvironmental influences*. J Immunol, 2005. **175**(1): p. 342-9.
28. Odegaard, J.I., et al., *Macrophage-specific PPARgamma controls alternative activation and improves insulin resistance*. Nature, 2007. **447**(7148): p. 1116-20.
29. Kang, K., et al., *Adipocyte-derived Th2 cytokines and myeloid PPARdelta regulate macrophage polarization and insulin sensitivity*. Cell Metab, 2008. **7**(6): p. 485-95.
30. Odegaard, J.I., et al., *Alternative M2 activation of Kupffer cells by PPARdelta ameliorates obesity-induced insulin resistance*. Cell Metab, 2008. **7**(6): p. 496-507.
31. Bouhrel, M.A., et al., *Unlike PPARgamma, PPARalpha or PPARbeta/delta activation does not promote human monocyte differentiation toward alternative macrophages*. Biochem Biophys Res Commun, 2009. **386**(3): p. 459-62.
32. Rubenfire, M., R.D. Brook, and R.S. Rosenson, *Treating mixed hyperlipidemia and the atherogenic lipid phenotype for prevention of cardiovascular events*. Am J Med, 2010. **123**(10): p. 892-8.
33. Drexel, H., *Statins, fibrates, nicotinic acid, cholesterol absorption inhibitors, anion-exchange resins, omega-3 fatty acids: which drugs for which patients?* Fundam Clin Pharmacol, 2009. **23**(6): p. 687-92.
34. Gordon, T., et al., *High density lipoprotein as a protective factor against coronary heart disease. The Framingham Study*. Am J Med, 1977. **62**(5): p. 707-14.
35. Fruchart, J.C., et al., *The Residual Risk Reduction Initiative: a call to action to reduce residual vascular risk in dyslipidaemic patient*. Diab Vasc Dis Res, 2008. **5**(4): p. 319-35.
36. Parhofer, K.G., *Beyond LDL-cholesterol: HDL-cholesterol as a target for atherosclerosis prevention*. Exp Clin Endocrinol Diabetes, 2005. **113**(8): p. 414-7.
37. Chinetti-Gbaguidi, G., J.C. Fruchart, and B. Staels, *Pleiotropic effects of fibrates*. Curr Atheroscler Rep, 2005. **7**(5): p. 396-401.
38. Fievet, C. and B. Staels, *Combination therapy of statins and fibrates in the management of cardiovascular risk*. Curr Opin Lipidol, 2009. **20**(6): p. 505-11.
39. Derosa, G. and P. Maffioli, *Effects of thiazolidinediones and sulfonylureas in patients with diabetes*. Diabetes Technol Ther, 2010. **12**(6): p. 491-501.
40. Calvert, J.W., et al., *Acute metformin therapy confers cardioprotection against myocardial infarction via AMPK-eNOS-mediated signaling*. Diabetes, 2008. **57**(3): p. 696-705.
41. Boden, G., et al., *Thiazolidinediones upregulate fatty acid uptake and oxidation in adipose tissue of diabetic patients*. Diabetes, 2005. **54**(3): p. 880-5.
42. Giri, S., K. Nieber, and A. Bader, *Hepatotoxicity and hepatic metabolism of available drugs: current problems and possible solutions in preclinical stages*. Expert Opin Drug Metab Toxicol, 2010. **6**(8): p. 895-917.
43. Karmen, A., F. Wroblewski, and J.S. Ladue, *Transaminase activity in human blood*. J Clin Invest, 1955. **34**(1): p. 126-31.
44. Lindblom, P., et al., *Isoforms of alanine aminotransferases in human tissues and serum--differential tissue expression using novel antibodies*. Arch Biochem Biophys, 2007. **466**(1): p. 66-77.
45. Nathwani, R.A., et al., *Serum alanine aminotransferase in skeletal muscle diseases*. Hepatology, 2005. **41**(2): p. 380-2.

46. Edgar, A.D., et al., *Fenofibrate modifies transaminase gene expression via a peroxisome proliferator activated receptor alpha-dependent pathway*. Toxicol Lett, 1998. **98**(1-2): p. 13-23.
47. Ozer, J., et al., *The current state of serum biomarkers of hepatotoxicity*. Toxicology, 2008. **245**(3): p. 194-205.
48. Lindena, J., et al., *Catalytic enzyme activity concentration in tissues of man, dog, rabbit, guinea pig, rat and mouse. Approach to a quantitative diagnostic enzymology, III. Communication*. J Clin Chem Clin Biochem, 1986. **24**(1): p. 35-47.
49. medicalbiochemistrypage.org.
50. Balfour, J.A., D. McTavish, and R.C. Heel, *Fenofibrate. A review of its pharmacodynamic and pharmacokinetic properties and therapeutic use in dyslipidaemia*. Drugs, 1990. **40**(2): p. 260-90.
51. Blane, G.F., *Comparative toxicity and safety profile of fenofibrate and other fibric acid derivatives*. Am J Med, 1987. **83**(5B): p. 26-36.
52. Bookout, A.L., et al., *Anatomical profiling of nuclear receptor expression reveals a hierarchical transcriptional network*. Cell, 2006. **126**(4): p. 789-99.
53. *A unified nomenclature system for the nuclear receptor superfamily*. Cell, 1999. **97**(2): p. 161-3.
54. Chawla, A., et al., *Nuclear receptors and lipid physiology: opening the X-files*. Science, 2001. **294**(5548): p. 1866-70.
55. Issemann, I. and S. Green, *Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators*. Nature, 1990. **347**(6294): p. 645-50.
56. Dreyer, C., et al., *Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors*. Cell, 1992. **68**(5): p. 879-87.
57. Kliewer, S.A., et al., *Differential expression and activation of a family of murine peroxisome proliferator-activated receptors*. Proc Natl Acad Sci U S A, 1994. **91**(15): p. 7355-9.
58. Ehrenborg, E. and A. Krook, *Regulation of skeletal muscle physiology and metabolism by peroxisome proliferator-activated receptor delta*. Pharmacol Rev, 2009. **61**(3): p. 373-93.
59. Kumar, R. and E.B. Thompson, *The structure of the nuclear hormone receptors*. Steroids, 1999. **64**(5): p. 310-9.
60. Gronemeyer, H., J.A. Gustafsson, and V. Laudet, *Principles for modulation of the nuclear receptor superfamily*. Nat Rev Drug Discov, 2004. **3**(11): p. 950-64.
61. Viswakarma, N., et al., *Coactivators in PPAR-Regulated Gene Expression*. PPAR Res, 2010. **2010**.
62. Spencer, T.E., et al., *Steroid receptor coactivator-1 is a histone acetyltransferase*. Nature, 1997. **389**(6647): p. 194-8.
63. Dowell, P., et al., *p300 functions as a coactivator for the peroxisome proliferator-activated receptor alpha*. J Biol Chem, 1997. **272**(52): p. 33435-43.
64. Zhu, Y., et al., *Cloning and identification of mouse steroid receptor coactivator-1 (mSRC-1), as a coactivator of peroxisome proliferator-activated receptor gamma*. Gene Expr, 1996. **6**(3): p. 185-95.
65. Nolte, R.T., et al., *Ligand binding and co-activator assembly of the peroxisome proliferator-activated receptor-gamma*. Nature, 1998. **395**(6698): p. 137-43.
66. Horlein, A.J., et al., *Ligand-independent repression by the thyroid hormone receptor mediated by a nuclear receptor co-repressor*. Nature, 1995. **377**(6548): p. 397-404.
67. Chen, J.D. and R.M. Evans, *A transcriptional co-repressor that interacts with nuclear hormone receptors*. Nature, 1995. **377**(6548): p. 454-7.
68. Yu, S. and J.K. Reddy, *Transcription coactivators for peroxisome proliferator-activated receptors*. Biochim Biophys Acta, 2007. **1771**(8): p. 936-51.
69. Dowell, P., et al., *Identification of nuclear receptor corepressor as a peroxisome proliferator-activated receptor alpha interacting protein*. J Biol Chem, 1999. **274**(22): p. 15901-7.

70. Zamir, I., J. Zhang, and M.A. Lazar, *Stoichiometric and steric principles governing repression by nuclear hormone receptors*. Genes Dev, 1997. **11**(7): p. 835-46.
71. Krogdham, A.M., et al., *Nuclear receptor corepressor-dependent repression of peroxisome-proliferator-activated receptor delta-mediated transactivation*. Biochem J, 2002. **363**(Pt 1): p. 157-65.
72. Shi, Y., M. Hon, and R.M. Evans, *The peroxisome proliferator-activated receptor delta, an integrator of transcriptional repression and nuclear receptor signaling*. Proc Natl Acad Sci U S A, 2002. **99**(5): p. 2613-8.
73. Lee, C.H., et al., *Transcriptional repression of atherogenic inflammation: modulation by PPARdelta*. Science, 2003. **302**(5644): p. 453-7.
74. Higgins, L.S. and A.M. Depaoli, *Selective peroxisome proliferator-activated receptor gamma (PPARgamma) modulation as a strategy for safer therapeutic PPARgamma activation*. Am J Clin Nutr, 2010. **91**(1): p. 267S-272S.
75. Barbier, O., et al., *Pleiotropic actions of peroxisome proliferator-activated receptors in lipid metabolism and atherosclerosis*. Arterioscler Thromb Vasc Biol, 2002. **22**(5): p. 717-26.
76. Juge-Aubry, C., et al., *DNA binding properties of peroxisome proliferator-activated receptor subtypes on various natural peroxisome proliferator response elements. Importance of the 5'-flanking region*. J Biol Chem, 1997. **272**(40): p. 25252-9.
77. Sher, T., et al., *cDNA cloning, chromosomal mapping, and functional characterization of the human peroxisome proliferator activated receptor*. Biochemistry, 1993. **32**(21): p. 5598-604.
78. Auboeuf, D., et al., *Tissue distribution and quantification of the expression of mRNAs of peroxisome proliferator-activated receptors and liver X receptor-alpha in humans: no alteration in adipose tissue of obese and NIDDM patients*. Diabetes, 1997. **46**(8): p. 1319-27.
79. Abbott, B.D., *Review of the expression of peroxisome proliferator-activated receptors alpha (PPAR alpha), beta (PPAR beta), and gamma (PPAR gamma) in rodent and human development*. Reprod Toxicol, 2009. **27**(3-4): p. 246-57.
80. Lefebvre, P., et al., *Sorting out the roles of PPAR alpha in energy metabolism and vascular homeostasis*. J Clin Invest, 2006. **116**(3): p. 571-80.
81. Kersten, S., et al., *Peroxisome proliferator-activated receptor alpha mediates the adaptive response to fasting*. J Clin Invest, 1999. **103**(11): p. 1489-98.
82. Kersten, S., et al., *The peroxisome proliferator-activated receptor alpha regulates amino acid metabolism*. Faseb J, 2001. **15**(11): p. 1971-8.
83. Staels, B., et al., *Fibrates downregulate apolipoprotein C-III expression independent of induction of peroxisomal acyl coenzyme A oxidase. A potential mechanism for the hypolipidemic action of fibrates*. J Clin Invest, 1995. **95**(2): p. 705-12.
84. Vu-Dac, N., et al., *Fibrates increase human apolipoprotein A-II expression through activation of the peroxisome proliferator-activated receptor*. J Clin Invest, 1995. **96**(2): p. 741-50.
85. Staels, B. and J. Auwerx, *Regulation of apo A-I gene expression by fibrates*. Atherosclerosis, 1998. **137 Suppl**: p. S19-23.
86. Delerive, P., et al., *Peroxisome proliferator-activated receptor alpha negatively regulates the vascular inflammatory gene response by negative cross-talk with transcription factors NF-kappaB and AP-1*. J Biol Chem, 1999. **274**(45): p. 32048-54.
87. Delerive, P., et al., *DNA binding-independent induction of IkappaBalpha gene transcription by PPARalpha*. Mol Endocrinol, 2002. **16**(5): p. 1029-39.
88. Madej, A., et al., *Effects of fenofibrate on plasma cytokine concentrations in patients with atherosclerosis and hyperlipoproteinemia IIb*. Int J Clin Pharmacol Ther, 1998. **36**(6): p. 345-9.
89. Fajas, L., et al., *The organization, promoter analysis, and expression of the human PPARgamma gene*. J Biol Chem, 1997. **272**(30): p. 18779-89.
90. Kepez, A., A. Oto, and S. Dagdelen, *Peroxisome proliferator-activated receptor-gamma: novel therapeutic target linking adiposity, insulin resistance, and atherosclerosis*. BioDrugs, 2006. **20**(2): p. 121-35.

91. Bugge, A. and S. Mandrup, *Molecular Mechanisms and Genome-Wide Aspects of PPAR Subtype Specific Transactivation*. PPAR Res, 2010. **2010**.
92. Skogsberg, J., et al., *Characterization of the human peroxisome proliferator activated receptor delta gene and its expression*. Int J Mol Med, 2000. **6**(1): p. 73-81.
93. Evans, R.M., G.D. Barish, and Y.X. Wang, *PPARs and the complex journey to obesity*. Nat Med, 2004. **10**(4): p. 355-61.
94. Tanaka, T., et al., *Activation of peroxisome proliferator-activated receptor delta induces fatty acid beta-oxidation in skeletal muscle and attenuates metabolic syndrome*. Proc Natl Acad Sci U S A, 2003. **100**(26): p. 15924-9.
95. Oliver, W.R., Jr., et al., *A selective peroxisome proliferator-activated receptor delta agonist promotes reverse cholesterol transport*. Proc Natl Acad Sci U S A, 2001. **98**(9): p. 5306-11.
96. Wang, Y.X., et al., *Peroxisome-proliferator-activated receptor delta activates fat metabolism to prevent obesity*. Cell, 2003. **113**(2): p. 159-70.
97. Luquet, S., et al., *Peroxisome proliferator-activated receptor delta controls muscle development and oxidative capability*. Faseb J, 2003. **17**(15): p. 2299-301.
98. Wang, Y.X., et al., *Regulation of muscle fiber type and running endurance by PPARdelta*. PLoS Biol, 2004. **2**(10): p. e294.
99. Barak, Y., et al., *Effects of peroxisome proliferator-activated receptor delta on placentation, adiposity, and colorectal cancer*. Proc Natl Acad Sci U S A, 2002. **99**(1): p. 303-8.
100. Peters, J.M., et al., *Growth, adipose, brain, and skin alterations resulting from targeted disruption of the mouse peroxisome proliferator-activated receptor beta(delta)*. Mol Cell Biol, 2000. **20**(14): p. 5119-28.
101. Desvergne, B. and W. Wahli, *Peroxisome proliferator-activated receptors: nuclear control of metabolism*. Endocr Rev, 1999. **20**(5): p. 649-88.
102. Xu, H.E., et al., *Structural determinants of ligand binding selectivity between the peroxisome proliferator-activated receptors*. Proc Natl Acad Sci U S A, 2001. **98**(24): p. 13919-24.
103. Johnson, T.E., et al., *Structural requirements and cell-type specificity for ligand activation of peroxisome proliferator-activated receptors*. J Steroid Biochem Mol Biol, 1997. **63**(1-3): p. 1-8.
104. Tan, N.S., et al., *Selective cooperation between fatty acid binding proteins and peroxisome proliferator-activated receptors in regulating transcription*. Mol Cell Biol, 2002. **22**(14): p. 5114-27.
105. Schug, T.T., et al., *Opposing effects of retinoic acid on cell growth result from alternate activation of two different nuclear receptors*. Cell, 2007. **129**(4): p. 723-33.
106. Brown, W.V., *Expert commentary: the safety of fibrates in lipid-lowering therapy*. Am J Cardiol, 2007. **99**(6A): p. 19C-21C.
107. Hodel, C., *Myopathy and rhabdomyolysis with lipid-lowering drugs*. Toxicol Lett, 2002. **128**(1-3): p. 159-68.
108. Boden, G. and M. Zhang, *Recent findings concerning thiazolidinediones in the treatment of diabetes*. Expert Opin Investig Drugs, 2006. **15**(3): p. 243-50.
109. Karalliedde, J. and R.E. Buckingham, *Thiazolidinediones and their fluid-related adverse effects: facts, fiction and putative management strategies*. Drug Saf, 2007. **30**(9): p. 741-53.
110. Grey, A., *Skeletal consequences of thiazolidinedione therapy*. Osteoporos Int, 2008. **19**(2): p. 129-37.
111. Henney, J.E., *From the Food and Drug Administration*. Jama, 2000. **283**(9): p. 1131.
112. Nissen, S.E. and K. Wolski, *Rosiglitazone Revisited: An Updated Meta-analysis of Risk for Myocardial Infarction and Cardiovascular Mortality*. Arch Intern Med, 2010.
113. Graham, D.J., et al., *Risk of acute myocardial infarction, stroke, heart failure, and death in elderly Medicare patients treated with rosiglitazone or pioglitazone*. Jama, 2010. **304**(4): p. 411-8.

114. Sprecher, D.L., et al., *Triglyceride:high-density lipoprotein cholesterol effects in healthy subjects administered a peroxisome proliferator activated receptor delta agonist*. *Arterioscler Thromb Vasc Biol*, 2007. **27**(2): p. 359-65.
115. Riserus, U., et al., *Activation of peroxisome proliferator-activated receptor (PPAR)delta promotes reversal of multiple metabolic abnormalities, reduces oxidative stress, and increases fatty acid oxidation in moderately obese men*. *Diabetes*, 2008. **57**(2): p. 332-9.
116. GlaxoSmithKline, *Protocol Summary for PAD100964* <http://www.gsk-clinicalstudyregister.com/index.jsp>. 2005.
117. Metabolex, <http://www.metabolex.com/news/nov182008.html>. 2008.
118. Feldman, P.L., M.H. Lambert, and B.R. Henke, *PPAR modulators and PPAR pan agonists for metabolic diseases: the next generation of drugs targeting peroxisome proliferator-activated receptors?* *Curr Top Med Chem*, 2008. **8**(9): p. 728-49.
119. Lundell, K., et al., *Alternative splicing of human peroxisome proliferator-activated receptor delta (PPAR delta): effects on translation efficiency and trans-activation ability*. *BMC Mol Biol*, 2007. **8**: p. 70.
120. Cresci, S., et al., *A PPARalpha promoter variant impairs ERR-dependent transactivation and decreases mortality after acute coronary ischemia in patients with diabetes*. *PLoS One*, 2010. **5**(9): p. e12584.
121. Clarke, S.L., C.E. Robinson, and J.M. Gimble, *CAAT/enhancer binding proteins directly modulate transcription from the peroxisome proliferator-activated receptor gamma 2 promoter*. *Biochem Biophys Res Commun*, 1997. **240**(1): p. 99-103.
122. Fajas, L., et al., *Regulation of peroxisome proliferator-activated receptor gamma expression by adipocyte differentiation and determination factor 1/sterol regulatory element binding protein 1: implications for adipocyte differentiation and metabolism*. *Mol Cell Biol*, 1999. **19**(8): p. 5495-503.
123. Keren, H., G. Lev-Maor, and G. Ast, *Alternative splicing and evolution: diversification, exon definition and function*. *Nat Rev Genet*, 2010. **11**(5): p. 345-55.
124. Jurica, M.S. and M.J. Moore, *Pre-mRNA splicing: awash in a sea of proteins*. *Mol Cell*, 2003. **12**(1): p. 5-14.
125. Zhou, Z., et al., *Comprehensive proteomic analysis of the human spliceosome*. *Nature*, 2002. **419**(6903): p. 182-5.
126. Black, D.L., *Mechanisms of alternative pre-messenger RNA splicing*. *Annu Rev Biochem*, 2003. **72**: p. 291-336.
127. Zahler, A.M., et al., *SR proteins: a conserved family of pre-mRNA splicing factors*. *Genes Dev*, 1992. **6**(5): p. 837-47.
128. Sanford, J.R., J. Ellis, and J.F. Caceres, *Multiple roles of arginine/serine-rich splicing factors in RNA processing*. *Biochem Soc Trans*, 2005. **33**(Pt 3): p. 443-6.
129. Smith, C.W. and J. Valcarcel, *Alternative pre-mRNA splicing: the logic of combinatorial control*. *Trends Biochem Sci*, 2000. **25**(8): p. 381-8.
130. Sugnet, C.W., et al., *Transcriptome and genome conservation of alternative splicing events in humans and mice*. *Pac Symp Biocomput*, 2004: p. 66-77.
131. Ast, G., *How did alternative splicing evolve?* *Nat Rev Genet*, 2004. **5**(10): p. 773-82.
132. Oakley, R.H., M. Sar, and J.A. Cidlowski, *The human glucocorticoid receptor beta isoform. Expression, biochemical properties, and putative function*. *J Biol Chem*, 1996. **271**(16): p. 9550-9.
133. Ogawa, S., et al., *Molecular cloning and characterization of human estrogen receptor betacx: a potential inhibitor of estrogen action in human*. *Nucleic Acids Res*, 1998. **26**(15): p. 3505-12.
134. Ebihara, K., et al., *Intron retention generates a novel isoform of the murine vitamin D receptor that acts in a dominant negative way on the vitamin D signaling pathway*. *Mol Cell Biol*, 1996. **16**(7): p. 3393-400.
135. Sabatino, L., et al., *A novel peroxisome proliferator-activated receptor gamma isoform with dominant negative activity generated by alternative splicing*. *J Biol Chem*, 2005. **280**(28): p. 26517-25.

136. Gervois, P., et al., *A truncated human peroxisome proliferator-activated receptor alpha splice variant with dominant negative activity*. Mol Endocrinol, 1999. **13**(9): p. 1535-49.
137. Bartel, D.P., *MicroRNAs: genomics, biogenesis, mechanism, and function*. Cell, 2004. **116**(2): p. 281-97.
138. Alvarez-Garcia, I. and E.A. Miska, *MicroRNA functions in animal development and human disease*. Development, 2005. **132**(21): p. 4653-62.
139. Rovira, C., M.C. Guida, and A. Cayota, *MicroRNAs and other small silencing RNAs in cancer*. IUBMB Life, 2010.
140. Chekulaeva, M. and W. Filipowicz, *Mechanisms of miRNA-mediated post-transcriptional regulation in animal cells*. Curr Opin Cell Biol, 2009. **21**(3): p. 452-60.
141. Chang, T.C. and J.T. Mendell, *microRNAs in vertebrate physiology and human disease*. Annu Rev Genomics Hum Genet, 2007. **8**: p. 215-39.
142. Schwarz, D.S., et al., *Asymmetry in the assembly of the RNAi enzyme complex*. Cell, 2003. **115**(2): p. 199-208.
143. Khvorova, A., A. Reynolds, and S.D. Jayasena, *Functional siRNAs and miRNAs exhibit strand bias*. Cell, 2003. **115**(2): p. 209-16.
144. Orom, U.A., F.C. Nielsen, and A.H. Lund, *MicroRNA-10a binds the 5'UTR of ribosomal protein mRNAs and enhances their translation*. Mol Cell, 2008. **30**(4): p. 460-71.
145. Duursma, A.M., et al., *miR-148 targets human DNMT3b protein coding region*. Rna, 2008. **14**(5): p. 872-7.
146. Nielsen, A.F., J. Gloggnitzer, and J. Martinez, *MicroRNAs cross the line: the battle for mRNA stability enters the coding sequence*. Mol Cell, 2009. **35**(2): p. 139-40.
147. Forman, J.J. and H.A. Collier, *The code within the code: MicroRNAs target coding regions*. Cell Cycle, 2010. **9**(8).
148. Grimson, A., et al., *MicroRNA targeting specificity in mammals: determinants beyond seed pairing*. Mol Cell, 2007. **27**(1): p. 91-105.
149. Lewis, B.P., et al., *Prediction of mammalian microRNA targets*. Cell, 2003. **115**(7): p. 787-98.
150. Bartel, D.P., *MicroRNAs: target recognition and regulatory functions*. Cell, 2009. **136**(2): p. 215-33.
151. Shin, C., et al., *Expanding the microRNA targeting code: functional sites with centered pairing*. Mol Cell, 2010. **38**(6): p. 789-802.
152. Martinelli, R., et al., *miR-519d Overexpression Is Associated With Human Obesity*. Obesity (Silver Spring), 2010. **18**(11): p. 2170-6.
153. Lin, Q., et al., *A role of miR-27 in the regulation of adipogenesis*. Febs J, 2009. **276**(8): p. 2348-58.
154. Kim, S.Y., et al., *miR-27a is a negative regulator of adipocyte differentiation via suppressing PPARgamma expression*. Biochem Biophys Res Commun, 2010. **392**(3): p. 323-8.
155. Jennewein, C., et al., *MicroRNA-27b contributes to lipopolysaccharide-mediated peroxisome proliferator-activated receptor gamma (PPARgamma) mRNA destabilization*. J Biol Chem, 2010. **285**(16): p. 11846-53.
156. Karbiener, M., et al., *microRNA miR-27b impairs human adipocyte differentiation and targets PPARgamma*. Biochem Biophys Res Commun, 2009. **390**(2): p. 247-51.
157. Gelman, L., et al., *Kinase signaling cascades that modulate peroxisome proliferator-activated receptors*. Curr Opin Cell Biol, 2005. **17**(2): p. 216-22.
158. Burns, K.A. and J.P. Vanden Heuvel, *Modulation of PPAR activity via phosphorylation*. Biochim Biophys Acta, 2007. **1771**(8): p. 952-60.
159. Lazennec, G., et al., *Activation of peroxisome proliferator-activated receptors (PPARs) by their ligands and protein kinase A activators*. Mol Endocrinol, 2000. **14**(12): p. 1962-75.
160. Hansen, J.B., et al., *Peroxisome proliferator-activated receptor delta (PPARdelta)-mediated regulation of preadipocyte proliferation and gene expression is dependent on cAMP signaling*. J Biol Chem, 2001. **276**(5): p. 3175-82.

161. Hirotani, M., et al., *Stabilization of peroxisome proliferator-activated receptor alpha by the ligand*. Biochem Biophys Res Commun, 2001. **288**(1): p. 106-10.
162. Hauser, S., et al., *Degradation of the peroxisome proliferator-activated receptor gamma is linked to ligand-dependent activation*. J Biol Chem, 2000. **275**(24): p. 18527-33.
163. Rieck, M., et al., *Expression level and agonist-binding affect the turnover, ubiquitination and complex formation of peroxisome proliferator activated receptor beta*. Febs J, 2007. **274**(19): p. 5068-76.
164. Feige, J.N., et al., *From molecular action to physiological outputs: peroxisome proliferator-activated receptors are nuclear receptors at the crossroads of key cellular functions*. Prog Lipid Res, 2006. **45**(2): p. 120-59.
165. Gill, G., *SUMO and ubiquitin in the nucleus: different functions, similar mechanisms?* Genes Dev, 2004. **18**(17): p. 2046-59.
166. Pourcet, B., et al., *SUMOylation of human peroxisome proliferator-activated receptor alpha inhibits its trans-activity through the recruitment of the nuclear corepressor NCoR*. J Biol Chem, 2010. **285**(9): p. 5983-92.
167. Leuenberger, N., S. Pradervand, and W. Wahli, *Sumoylated PPARalpha mediates sex-specific gene repression and protects the liver from estrogen-induced toxicity in mice*. J Clin Invest, 2009. **119**(10): p. 3138-48.
168. Pascual, G., et al., *A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR-gamma*. Nature, 2005. **437**(7059): p. 759-63.
169. Glinghammar, B., et al., *PPARdelta activation induces COX-2 gene expression and cell proliferation in human hepatocellular carcinoma cells*. Biochem Biophys Res Commun, 2003. **308**(2): p. 361-8.
170. Cartharius, K., et al., *MatInspector and beyond: promoter analysis based on transcription factor binding sites*. Bioinformatics, 2005. **21**(13): p. 2933-42.
171. Grabe, N., *AliBaba2: context specific identification of transcription factor binding sites*. In Silico Biol, 2002. **2**(1): p. S1-15.
172. Messeguer, X., et al., *PROMO: detection of known transcription regulatory elements using species-tailored searches*. Bioinformatics, 2002. **18**(2): p. 333-4.
173. Ovcharenko, I., et al., *Mulan: multiple-sequence local alignment and visualization for studying function and evolution*. Genome Res, 2005. **15**(1): p. 184-94.
174. ClustalW, <http://www.ebi.ac.uk/clustalw>.
175. RepeatMasker, <http://www.repeatmasker.org>.
176. UCSCgenomebrowser, <http://genome.ucsc.edu/>.
177. TargetScanS, <http://genes.mit.edu/tscan/targetscanS2005.html>.
178. PicTar, <http://pictar.mdc-berlin.de/>.
179. Repnik, U., M. Knezevic, and M. Jeras, *Simple and cost-effective isolation of monocytes from buffy coats*. J Immunol Methods, 2003. **278**(1-2): p. 283-92.
180. Vosper, H., et al., *The peroxisome proliferator-activated receptor delta promotes lipid accumulation in human macrophages*. J Biol Chem, 2001. **276**(47): p. 44258-65.
181. Laemmli, U.K., *Cleavage of structural proteins during the assembly of the head of bacteriophage T4*. Nature, 1970. **227**(5259): p. 680-5.
182. Birjmohun, R.S., et al., *Apolipoprotein A-II is inversely associated with risk of future coronary artery disease*. Circulation, 2007. **116**(18): p. 2029-35.
183. Ladias, J.A., et al., *Transcriptional regulation of human apolipoprotein genes ApoB, ApoCIII, and ApoAII by members of the steroid hormone receptor superfamily HNF-4, ARP-1, EAR-2, and EAR-3*. J Biol Chem, 1992. **267**(22): p. 15849-60.
184. Ikewaki, K., et al., *Apolipoprotein A-II production rate is a major factor regulating the distribution of apolipoprotein A-I among HDL subclasses LpA-I and LpA-I:A-II in normolipidemic humans*. Arterioscler Thromb Vasc Biol, 1995. **15**(3): p. 306-12.
185. Silva, R.A., et al., *The structure of apolipoprotein A-II in discoidal high density lipoproteins*. J Biol Chem, 2007. **282**(13): p. 9713-21.

186. Tomkiewicz, C., et al., *Opposite regulation of the rat and human cytosolic aspartate aminotransferase genes by fibrates*. *Biochem Pharmacol*, 2004. **67**(2): p. 213-25.
187. Friis, R., et al., *Effect of rapid weight loss with supplemented fasting on liver tests*. *J Clin Gastroenterol*, 1987. **9**(2): p. 204-7.
188. Horio, Y., et al., *Rat cytosolic aspartate aminotransferase: regulation of its mRNA and contribution to gluconeogenesis*. *J Biochem*, 1988. **103**(5): p. 805-8.
189. Palou, A., et al., *Changes in alanine transaminase activity in several organs of the rat induced by a 24-hour fast*. *Horm Metab Res*, 1980. **12**(10): p. 505-8.
190. Sattar, N., et al., *Elevated alanine aminotransferase predicts new-onset type 2 diabetes independently of classical risk factors, metabolic syndrome, and C-reactive protein in the west of Scotland coronary prevention study*. *Diabetes*, 2004. **53**(11): p. 2855-60.
191. Hanley, A.J., et al., *Elevations in markers of liver injury and risk of type 2 diabetes: the insulin resistance atherosclerosis study*. *Diabetes*, 2004. **53**(10): p. 2623-32.
192. Schindhelm, R.K., et al., *Alanine aminotransferase predicts coronary heart disease events: a 10-year follow-up of the Hoorn Study*. *Atherosclerosis*, 2007. **191**(2): p. 391-6.
193. Schindhelm, R.K., et al., *Alanine aminotransferase as a marker of non-alcoholic fatty liver disease in relation to type 2 diabetes mellitus and cardiovascular disease*. *Diabetes Metab Res Rev*, 2006. **22**(6): p. 437-43.
194. Belcher, G. and G. Schernthaner, *Changes in liver tests during 1-year treatment of patients with Type 2 diabetes with pioglitazone, metformin or gliclazide*. *Diabet Med*, 2005. **22**(8): p. 973-9.
195. Gastaldelli, A., et al., *The effect of rosiglitazone on the liver: decreased gluconeogenesis in patients with type 2 diabetes*. *J Clin Endocrinol Metab*, 2006. **91**(3): p. 806-12.
196. Jackson, E.R., et al., *The early effects of short-term dexamethasone administration on hepatic and serum alanine aminotransferase in the rat*. *Drug Chem Toxicol*, 2008. **31**(4): p. 427-45.
197. Bastie, C., et al., *Alterations of peroxisome proliferator-activated receptor delta activity affect fatty acid-controlled adipose differentiation*. *J Biol Chem*, 2000. **275**(49): p. 38768-73.
198. Pesant, M., et al., *Peroxisome proliferator-activated receptor delta (PPARdelta) activation protects H9c2 cardiomyoblasts from oxidative stress-induced apoptosis*. *Cardiovasc Res*, 2006. **69**(2): p. 440-9.
199. Larsen, L.K., et al., *Genomic organization of the mouse peroxisome proliferator-activated receptor beta/delta gene: alternative promoter usage and splicing yield transcripts exhibiting differential translational efficiency*. *Biochem J*, 2002. **366**(Pt 3): p. 767-75.
200. Chew, C.H., et al., *Molecular characterisation of six alternatively spliced variants and a novel promoter in human peroxisome proliferator-activated receptor alpha*. *Biochem Biophys Res Commun*, 2003. **305**(2): p. 235-43.
201. Chen, Y., A.R. Jimenez, and J.D. Medh, *Identification and regulation of novel PPAR-gamma splice variants in human THP-1 macrophages*. *Biochim Biophys Acta*, 2006. **1759**(1-2): p. 32-43.
202. McClelland, S., R. Shrivastava, and J.D. Medh, *Regulation of Translational Efficiency by Disparate 5' UTRs of PPARgamma Splice Variants*. *PPAR Res*, 2009. **2009**: p. 193413.
203. Wang, X.Q. and J.A. Rothnagel, *5'-untranslated regions with multiple upstream AUG codons can support low-level translation via leaky scanning and reinitiation*. *Nucleic Acids Res*, 2004. **32**(4): p. 1382-91.
204. Bazzoni, F., et al., *Induction and regulatory function of miR-9 in human monocytes and neutrophils exposed to proinflammatory signals*. *Proc Natl Acad Sci U S A*, 2009. **106**(13): p. 5282-7.
205. Krichevsky, A.M., et al., *Specific microRNAs modulate embryonic stem cell-derived neurogenesis*. *Stem Cells*, 2006. **24**(4): p. 857-64.

206. Plaisance, V., et al., *MicroRNA-9 controls the expression of Granuphilin/Slp4 and the secretory response of insulin-producing cells*. J Biol Chem, 2006. **281**(37): p. 26932-42.
207. Barish, G.D., V.A. Narkar, and R.M. Evans, *PPAR delta: a dagger in the heart of the metabolic syndrome*. J Clin Invest, 2006. **116**(3): p. 590-7.
208. Narkar, V.A., et al., *AMPK and PPARdelta agonists are exercise mimetics*. Cell, 2008. **134**(3): p. 405-15.